

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/329979010>

First record of *Calycuoniscus goeldii* (Lemos de Castro, 1967) (Oniscidea: Dubioniscidae) for the state of Piauí, northeastern Brazil

Article in *Pesquisa e Ensino em Ciências Exatas e da Natureza* - December 2018

DOI: 10.29215/pecen.v2i2.1069

CITATIONS

0

READS

142

3 authors:



Vanderley Ferreira Silva Costa
Universidade Federal do Ceará

1 PUBLICATION 0 CITATIONS

[SEE PROFILE](#)



Daniela Grangeiro
Universidade Estadual do Piauí (UESPI)

6 PUBLICATIONS 15 CITATIONS

[SEE PROFILE](#)



Carlos Anderson Soares Bezerra Pereira
Universidade Federal do Piauí

1 PUBLICATION 0 CITATIONS

[SEE PROFILE](#)


Some of the authors of this publication are also working on these related projects:



Análise filogenética dos Prosekiini Leistikow, 2001 [View project](#)



Isopoda (CRUSTACEA: ONISCIDEA) from the State of Piauí [View project](#)



First record of *Calycuoniscus goeldii* (Lemos de Castro, 1967) (Oniscidea: Dubioniscidae) for the state of Piauí, northeastern Brazil

Vanderley Ferreira Silva Costa¹, Daniela Correia Grangeiro^{1,2} & Carlos Anderson Soares Bezerra Pereira¹

- (1) Universidade Estadual do Piauí – Campus Professor Barros Araújo, Av. Senador Helvídio Nunes de Barros Junco, Picos 64600000, Piauí, Brazil. E-mail: yderley@gmail.com
- (2) Universidade Federal da Paraíba – Campus I, Centro de Ciências Exatas e da Natureza, Departamento de Sistemática e Ecologia, Programa de Pós-Graduação em Ciências Biológicas (Zoologia), Cidade Universitária 58051-900, João Pessoa, Paraíba, Brazil. E-mail: daniela.grangeiro@gmail.com

Costa V.F.S., Grangeiro D.C. & Pereira C.A.S.B. (2018) First record of *Calycuoniscus goeldii* (Lemos de Castro, 1967) (Oniscidea: Dubioniscidae) for the state of Piauí, northeastern Brazil. *Pesquisa e Ensino em Ciências Exatas e da Natureza*, 2(2): 152–157. <http://dx.doi.org/10.29215/pecen.v2i2.1069>

Academic editor: Flavio A. Alves-Jr. **Received:** 23 October 2018. **Accepted:** 28 November 2018. **Published:** 28 December 2018.

Primeiro registro de *Calycuoniscus goeldii* (Lemos de Castro, 1967) (Oniscidea, Dubioniscidae) para o Estado do Piauí, Nordeste do Brasil

Resumo: O presente trabalho fornece o primeiro registro de *Calycuoniscus goeldii* (Lemos de Castro, 1967) para o município de Picos, estado do Piauí, nordeste do Brasil. Além disso, este trabalho contribui para ampliar o conhecimento sobre a biodiversidade de isópodes terrestres do estado do Piauí e distribuição da família Dubioniscidae no Nordeste do Brasil.

Palavras-chave: Crustacea, Isopoda, novo registro, Picos.

Abstract: The present work provides the first record of *Calycuoniscus goeldii* (Lemos de Castro, 1967) from the municipality of Picos, state of Piauí, northeastern Brazil. In addition, this paper contributes to increase the knowledge about the biodiversity of terrestrial isopods of the state of Piauí and distribution of the family Dubioniscidae in northeastern Brazil.

Key words: Crustacea, Isopoda, new record, Picos.

Introduction

Isopods of the suborder Oniscidea Latreille, 1802 include most terrestrial species (Schmidt 2008). This group consists of about 3.710 species belonging to 38 families and 527 genera (Schmalfuss 2003; Sfenthourakis & Taiti 2015). Most members of the suborder are small to medium in size, with species distributed worldwide from desert regions to forests, including habitats such as meadows, agroecosystems, mountains and caves (Hopkin & Martin 1984).

Members of the Oniscidea have a relevant ecological role in terrestrial ecosystems since they feed on organic matter contributing to the nutrient cycling deposited in the soil as fecal pellets and in the formation of humus (Quadros & Araújo 2008).

There are about 180 species of Recent terrestrial isopods in Brazil (Souza-Kury 1998; Schmalfuss 2003; Souza & Grangeiro 2006; Grangeiro *et al.* 2014; Campos-Filho *et al.* 2014, 2015;

Cardoso *et al.* 2016; Campos-Filho *et al.* 2018). Oniscidea from the family Dubioniscidae Schultz, 1995 are widely distributed in South America mainly in Venezuela, Trindade, Guyana and Brazil (between the northern to southeastern of the country), Paraguay and Argentina. Dubioniscidae includes the following genera: *Dubioniscus* Vandel, 1963; *Calycuoniscus* Collinge, 1915; and *Novamundoniscus* Schultz, 1995 (Lemos de Castro 1967, 1968; Souza-Kury 1998; Schmalfluss 2003; Campos-Filho *et al.* 2014; Grangeiro *et al.* 2014). The family is characterized by the presence of bristles on the dorsal region, cephalothorax with central-frontal depression and well-developed lateral lobes, antenna of three articles, pleon as wide as the pereon with well developed neopleurae, male and female with exopods of pleopods having similar shape (without respiratory structures) and exopods of the uropods surpassing the size of the pleotelson (Schultz 1995). Schmidt (2003) states that all characters attributed to Dubioniscidae are plesiomorphic or ambiguous and that the family may be paraphyletic.

The biological and ecological knowledge of Oniscidea from northeastern Brazil is still very incipient. The state of Piauí is one of the regions in the northeast of the country in which the group was little studied with only the records of the following families: Porcellionidae Brandt & Raterburg, 1831; Platyarthridae Verhoeff, 1949; Armadillidae Brandt & Raterburg, 1831; and Dubioniscidae Schultz, 1995 (Grangeiro *et al.* 2014).

This study provides the first record of *Calycuoniscus goeldii* (Lemos de Castro, 1967) for the state of Piauí, northeastern Brazil.

Material and Methods

Study area

The specimens examined in this study were collected between 2013 and 2017 in the vicinity of the Conduru and Junco neighborhood, both located in the municipality of Picos, state of Piauí, northeastern Brazil (Figure 1). The municipality is located in agricultural production systems of the state of Piauí, located 310 km from Teresina (Sousa 2012).




Figure 1. (A) Map of Brazil highlighting the state of Piauí; (B) Map of the city of Picos showing the neighborhoods of Conduru and Junco; (C) Collection area in the neighborhood of Junco (purple triangle – sampling site); (D) Collection area in the neighborhood of Conduru (green pentagon – sampling site).

Sampling, sorting and identification

The samples were carried out through active search by members of the *Grupo de Estudos Taxonômicos dos Oniscidea do Piauí* (GETOPI). Specimens were stored in ethanol 70% and deposited in the *Laboratório de Biologia da Universidade Estadual do Piauí* (UESPI), Campus Professor Barros Araújo, Picos, Piauí. All specimens were analyzed under stereomicroscope with the aid of brushes, tweezers and petri dishes. The identification was performed based on Lemos de Castro (1967, 1968) and Schultz (1995) with aid of stereomicroscope and optical microscope through the morphological analysis of the *habitus*, antenna and pereopods of individuals. The material analyzed is deposited and available for study at the “Universidade Estadual do Piauí (UESPI)”, Picos, Piauí and “Coleção de Invertebrados Paulo Young, Departamento de Sistemática e Ecologia, Universidade Federal da Paraíba (UFPB)”, João Pessoa, Paraíba, Brazil”.

Results

Isopoda Latreille, 1817
Oniscidea Latreille, 1803
Dubioniscidae Schultz, 1995
Calycuoniscus Collinge, 1915

Type species: *Calycuoniscus bodkini* Collinge, 1915 by monotypy.

Calycuoniscus goeldii (Lemos de Castro, 1967)
Figures 2–3

Hileioniscus goeldii Lemos de Castro 1967: 318.

Calycuoniscus goeldii – Lemos de Castro (1968: 409, figs 1–7); Cardoso *et al.* (2016: 3).

Dubioniscus goeldii – Schultz (1995: 400).

Material examined: [09♀] Brazil, Piauí, Picos, Condurú, 07°04'91" S, 41°25'47" W, 12.VII.2013, collector GETOPI (UESPI-LB 0016); [1♂] Brazil, Piauí, Picos, Condurú, 07°04'91" S, 41°25'47" W, 12.VII.2013, collector GETOPI (UESPI-LB 0018); [1 ♀] Brazil, Piauí, Picos, Condurú, 07°04'91" S, 41°25'47" W, 26-IX-2015, collector GETOPI (UESPI-LB 0071); [2♀] Brazil, Piauí, Picos, Condurú, 07°04'91" S, 41°25'47" W, 30.X.2015, collector GETOPI (UESPI-LB 0091); [5♀] Brazil, Piauí, Picos, Junco, 07°04'44" S, 41°26'33" W, 02.X.2016, collector GETOPI (UESPI-LB 0093); [2♀] Brazil, Piauí, Picos, Condurú, 07°04'91" S, 41°25'47" W, 07.III.2017, collector GETOPI (UESPI-LB 0100); [02♀] Brazil, Piauí, Picos, Condurú, 07°04'91" S, 41°25'47" W, 12.VII.2013, collector GETOPI (UFPB).

Diagnosis: Specimens having 5 mm in maximum length. Brown color with light spots and continuous light longitudinal bands between median line and lateral margins of pereonites and pleon. Tegument covered by scaly setae. Cephalothorax without frontal depression, with triangular-shaped medial and lateral lobes with straight sides. Antenna having three articles with thin suture between second and third. Male with exopod of pleopode 1 oval and widened distal part; endopod with distal end having row of small spines. Male with exopod of pleopode 2 triangular. Uropod exopods surpassing telson; endopods not visible in dorsal view.

Geographic distribution: BRAZIL – Pará (Lemos de Castro 1967, 1968; Souza-Kury 1998; Leistikow & Wägele 1999; Schmalfuss 2003), Piauí (Picos) (present study), Chapada do Araripe (Crato-Ceará) (Souza & Grangeiro 2006) (Figure 1).

Remarks: The individuals of *Calycuoniscus goeldii* were recognized here mainly based on antenna with thin suture between the second and third articles (Figure 3), bristles of *habitus*

directed backwards, brown color with light spots (Figure 2) and cephalothorax with median and lateral lobes with triangular shape and straight sides.



Figure 2. Habitus of female individual of *Calycuoniscus goeldii* collected in the state of Piauí (Picos), northeastern Brazil.



Figure 3. Antenna with thin suture between the second and third articles of female individual of *Calycuoniscus goeldii* collected in the state of Piauí (Picos), northeastern Brazil.

Discussion

Members of the genera *Dubioniscus* and *Calycuoniscus* are similar in general morphology and antenna with thin suture between the second and third articles in an approximately bi-flagellated condition. Furthermore, *Dubioniscus* has a V-shape anterodorsal depression in the cephalothorax (Schultz 1995), which is absent in individuals of the genera *Calycuoniscus* and *Novamundoniscus* (Cardoso *et al.* 2016).

To date, only four species of the genus *Calycuoniscus* are known: *C. ambiguus* (Budde-Lund, 1893), *C. compar* (Budde-Lund, 1893) [Venezuela], *C. bodkini* Collinge, 1915 [Trindade, Guiana and Brazil] and *C. spinosus* Collinge, 1917 [Trindade (West Indies)] (Schultz 1995). There are now two species of this genus with occurrences in Brazil: *C. bodkini* Collinge, 1915 recorded from Amazon rainforest (Schmalfuss 2003) and *C. goeldii* Lemos de Castro, 1967 recorded from the states of Pará and Ceará (Schmalfuss 2003; Souza & Grangeiro 2006).

Calycuoniscus goeldii is recorded from cascade region of the Batateiras river, Chapada do Araripe, state of Ceará, northeastern Brazil (Souza & Grangeiro 2006). This study represents the first record of *C. goeldii* from the state of Piauí and it contributes to expand the knowledge on the species richness in the region and the geographic distribution of taxa belonging to the family Dubioniscidae in northeastern Brazil.

Acknowledgments

The authors are grateful to M. Sc. Rudá Amorim Lucena (UFPB/CCEN/DSE) and the anonymous reviewers for their precious comments on the paper.

References

- Campos-Filho I.S., Araujo P.B., Bichuette M.E., Trajano E. & Taiti S. (2014) Terrestrial isopods (Crustacea: Isopoda: Oniscidea) from Brazilian caves. *Zoological Journal of the Linnean Society*, 172(2): 360–425. DOI: 10.1111/zoj.12172
- Campos-Filho I.S., Mise K.M. & Sessegolo G.C. (2015) A new species of *Trichorhina* Budde-Lund, 1908 (Isopoda: Oniscidea: Platyarthridae) from Paraná caves, southern Brazil. *Nauplius*, 23(2): 112–119. DOI: 10.1590/S0104-64972015002324
- Campos-Filho I.S., Lisboa J.T. & Cardoso G.M. (2018) A new genus and two new species of Pudeoniscidae Lemos de Castro 1973 (Crustacea: Isopoda: Oniscidea) from Brazil. *Journal of Natural History*, 52(7-8): 457–482, DOI: 10.1080/00222933.2018.1437229.
- Cardoso G.M., Campos-Filho I.S. & Araujo P.B. (2016) The genus *Dubioniscus* Vandel, 1963 (Oniscidea, Dubioniscidae) with descriptions of two new species from Brazil. *Tropical Zoology*, 29: 111–133. DOI: 10.1080/03946975.2016.1179024
- Grangeiro D.C., Borges N.R.S. & Moura G.I.B.S.C. (2014) Primeiro registro de Crustáceos Terrestres (Isopoda, Oniscidea) para o Piauí, Brasil. (p. 89–101). *In*: Grangeiro D.C., Azar G.S. & Pessoa W.R.L.S. (Orgs). Pesquisas no Semiárido Piauiense. Curitiba: CRV. 204 p.
- Hopkin S.P. & Martin M.H. (1984) Heavy metals in woodlice (p. 143–166). *In*: Sutton S.L. & Holdich D.M. (Eds). The Biology of Terrestrial Isopods. The Zoological Society of London. Number 53. Oxford: Clarendon Press. 518 p.
- Leistikow A. & Wägele J. (1999) Checklist of the terrestrial isopods of the new world (Crustacea, Isopoda, Oniscidea). *Revista Brasileira de Zoologia*, 16: 1–72. DOI: 10.1590/S0101-81751999000100001
- Lemos de Castro A. (1967) Isópodos terrestres da Amazônia Brasileira (Isopoda, Oniscoidea). *Atas do Simpósio sobre a Biota Amazônica*, 5: 311–336.
- Lemos de Castro A. (1968) Descrição complementar de *Calycuoniscus goeldii* (Lemos de Castro) (Isopoda Terrestria, Oniscidae, Bathytropinae). *Revista Brasileira de Biologia*, 28(4): 407–412.
- Quadros A.F. & Araújo P.B. (2008) An assemblage of terrestrial isopods (Crustacea) in southern Brazil and its contribution to leaf litter processing. *Revista Brasileira de Zoologia* 25(1): 58–66. DOI: 10.1590/S0101-81752008000100009
- Schmalzfuss H. (2003) World catalog of terrestrial isopods (Isopoda: Oniscidea). *Stuttgater Beiträge zur Naturkunde*, 654: 1–341.
- Schmidt C. (2003) Contribution to the phylogenetic system of the Crinocheta (Crustacea, Isopoda). Part 2. (Oniscoidea to Armadillidiidae). *Mitteilungen aus dem Museum für Naturkunde in Berlin, Zoologische Reihe*, 79: 3–179. DOI: 10.1002/mmzn.4850790102
- Schmidt C. (2008) Phylogeny of the Terrestrial Isopoda (Oniscidea): a Review. *Arthropod Systematics & Phylogeny*, 66(2): 191–226.
- Schultz G.A. (1995) Terrestrial isopod crustaceans (Oniscidea) from Paraguay with definition of a new family. *Revue Suisse de Zoologie*, 102: 387–424.
- Sfenthourakis S. & Taiti S. (2015) Patterns of taxonomic diversity among terrestrial isopods. *ZooKeys*, 515: 13–25. DOI: 10.3897/zookeys.515.9332
- Sousa M.D. (2012) Identidade e docência: o saber-fazer do professor e de Sociologia das escolas públicas estaduais de Picos/PI. Tese, Programa de Pós-Graduação em Educação, Universidade Federal do Rio Grande do Norte, Natal.
- Souza L.A. & Grangeiro D.C. (2006) Primeiro registro de crustáceos terrestres (Isopoda, Oniscidea) para a Chapada do Araripe, Ceará, Brasil. *Cadernos de Cultura e Ciência*, 1(1): 33–39.

Calycuoniscus goeldii for the state of Piauí

Souza-Kury L.A. (1998) Malacostraca - Peracarida. Isopoda. Oniscidea (p. 653–674). *In*: Young P.S. (Ed.). Catalogue of Crustacea of Brazil. Rio de Janeiro: Museu Nacional. 717 p.