

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/343658364>

# Micropredator Behaviour of *Rocinela dumerilii* (Isopoda, Aegidae) on *Trachurus trachurus* in the Sea of Marmara (Turkey)

Article in *Transylvanian Review of Systematical and Ecological Research* · August 2020

DOI: 10.2478/trser-2020-0011

CITATIONS

0

READS

307

3 authors, including:



Ahmet Öktener

135 PUBLICATIONS 780 CITATIONS

SEE PROFILE



Murat Şirin

Republic of Turkey Ministry of Agriculture and Forestry

14 PUBLICATIONS 6 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



TAGEM/HAYSUD/2014/05/01 [View project](#)



AGEM/HAYSUD/2011/A11/P-02/2 [View project](#)

**MICROPREDATOR BEHAVIOUR  
OF *ROCINELA DUMERILII* (ISOPODA, AEGIDAE)  
ON *TRACHURUS TRACHURUS*  
IN THE SEA OF MARMARA (TURKEY)**

*Ahmet ÖKTENER* \*, *Murat ŞİRİN* \* and *Emrah YURDİGÜL* \*

\* Sheep Research Institute, Department of Fisheries, Çanakkele Road 7 km, Bandırma, Balıkesir, Turkey, TR-10200, ahmetoktener@yahoo.com, mothocya@hotmail.com, emrah.yurdigul@tarim.gov.tr

DOI: 10.2478/trser-2020-0011

**KEYWORDS:** *Rocinela dumerilii*, micropredator, *Trachurus trachurus*, *Elthusa sinuata*, the Sea of Marmara.

**ABSTRACT**

*Rocinela dumerilii* (Lucas, 1849) is obtained on the horse mackerel, *Trachurus trachurus* (Linnaeus, 1758) from the Sea of Marmara, in Turkey. This species is a free-living organism from the Aegean Sea of Turkey, the Mediterranean Sea and Atlantic. Its micropredator behaviour is discovered for the first time in this study. *Elthusa sinuata* (Koelbel, 1879) is also collected on the gill cavity of Fries's goby, *Lesueurigobius friesii* (Malm, 1874) in this study. This host is a new record for *Elthusa sinuata*. The morphological characters of *Rocinela dumerilii* species are given by drawings.

**RÉSUMÉ:** Le comportement microprédateur de *Rocinela dumerilii* (Isopoda, Aegidae) sur le chinchard commun *Trachurus trachurus* dans la Mer de Marmara (Turquie).

*Rocinela dumerilii* (Lucas, 1849) est collectée sur les exemplaires de chinchard *Trachurus trachurus* (Linnaeus, 1758) en provenant de la Mer de Marmara, en Turquie. Cette espèce est connue en tant que forme de vie libre, de la Mer Egée, en Turquie, de la Mer Méditerranée et de l'Atlantique. Son comportement micro-prédateur a été découvert pour la première fois durant cette étude. *Elthusa sinuata* (Koelbel, 1879) est également collecté de la cavité branchiale du gobie à grandes écailles, *Lesueurigobius friesii* (Malm, 1874) durant cette étude. Cet hôte est un nouveau record pour *Elthusa sinuata*. Les caractères morphologiques de l'espèce de *Rocinela dumerilii* sont indiqués par des dessins.

**REZUMAT:** Comportamentul microprădător al speciei *Rocinela dumerilii* (Isopoda, Aegidae) pe stavridul atlantic *Trachurus trachurus* în Marea Marmara (Turcia).

*Rocinela dumerilii* (Lucas, 1849) este colectată de pe exemplare de stavrid atlantic *Trachurus trachurus* (Linnaeus, 1758) din Marea Marmara, Turcia. Această specie este cunoscută ca formă de viață liberă, în Marea Egee, în Turcia, din Marea Mediterană și Atlantic. Comportamentul său microprădător a fost descoperit pentru prima dată în timpul acestui studiu. *Elthusa sinuata* (Koelbel, 1879) a fost, de asemenea, colectată din cavitatea branhială a guvidelui lui Fries, *Lesueurigobius friesii* (Malm, 1874) în timpul acestui studiu. Această gazdă este o nouă semnalare pentru *Elthusa sinuata*. Caracterile morfologice ale speciei *Rocinela dumerilii* sunt indicate prin desene.

## INTRODUCTION

Members of the family Aegidae are active predators on hosts such as fish and elasmobranches (Brusca, 1983; Bruce, 2003, 2004). These isopods attack the external surfaces of fish. They were also reported as associated with invertebrates (Brusca, 1983; Bruce, 2003, 2004). They may be named as micropredators or temporary parasites. They may also act as scavengers, consuming dead fish that settle to the benthos. When not feeding, aegids lie on the substratum (Brusca and Iverson, 1985). Aegidae are distributed throughout the world oceans, from the tropics to polar waters (Bruce, 2009). Forty-one species of *Rocinela* are listed by Worms Editorial Board (2019).

There are few association records between aegids, fish, and invertebrates hosts: *Rocinela signata* was reported on the general body surface and gills of hosts such as nurse sharks (Brusca and Iverson, 1985). *R. signata* was reported from *Dasyatis americana*, *D. guttata* (Williams et al., 1994); from few fish species including *Archosargus probatocephalus*, *Balistes vetula*, *Bothus lunatus*, *Calamus calamus*, *Calamus bajonado*, *Calamus penna*, *Caranx* sp., *Dasyatis americana*, *Epinephelus itajara*, *Epinephelus morio*, *Galeocerdo cuvieri*, *Ginglymostoma cirratum*, *Haemulon steindachneri*, *Haemulon flavolineatum*, *Lachnolaimus maximus*, *Lutjanus analis*, *Lutjanus buccanella*, *Lutjanus blackfordi*, *Mycteroperca venenosa*, *Mycteroperca bonaci*, *Orthopristis ruber*, *Raja eglanteria*, *Sparisoma viride*, *Sphyrna barracuda* (Kensley and Schotte, 1989); from *Lutjanus analis*, *Micropogonias furnieri* (Bunkley-Williams and Williams, 1998). *Aega tridens* was reported from *Scomber scomber* (Treasurer, 2001); while *Aega rosacea* from *Epinephelus guaza*, *Mullus barbatus barbatus*, *Squatina squatina* and *Aega deshaysiana* from *Sardina pilchardus* (Ramdane and Trilles, 2008). Garzón-Ferreira (1990) also noted *R. signata* attacking humans in Colombia.

The *Elthusa* members are observed in the branchial cavities of hosts. 35 species belonging to this genus are listed by Worms Editorial Board (2019). It is recorded from the Indo-West Pacific, Eastern Pacific, the Atlantic, and the Mediterranean (van der Wal et al., 2019). *Elthusa sinuata* (Koelbel, 1879) is known from the Mediterranean Sea red bandfish.

Although species of *Rocinela* from Mediterranean has been reported by several researchers many times, the morphology of females and males of this genus in the Mediterranean has not been investigated. Specific key is necessary to identify the species belonging *Rocinela* genus for Mediterranean Sea.

The aim of this study is to provide a tool for taxonomists for future use. The present study contributes the presence of micropredator behavior of *R. dumerilii* on host fish and also to the knowledge of the host diversity of *Elthusa sinuata*.

## MATERIAL AND METHODS

*Trachurus trachurus* (Linnaeus, 1758) (Fig. 1) (Perciformes, Carangidae) and *Lesueurigobius friesii* (Malm, 1874) (Fig. 2) (Perciformes, Gobiidae) were caught by trawling net in Marmara Sea, Turkey, on July 2017. Collected isopod samples were fixed in 70% ethanol. Some specimens were dissected with the use of a Wild M5 stereo microscope. The appendages were drawn with the aid of a camera lucida (Olympus BH-DA). Drawings were transferred to digital media via scanner. The photos were taken with a Canon EOS 1100D camera attached to a microscope. Measurements (in  $\mu\text{m}$ ) were obtained using a program (Pro-way). Identifications and comparisons were performed according to Schiodte and Meinert (1879), Kussakin (1979), Monod (1923) for *R. dumerilii* and Trilles (1976), Ökterner et al. (2018) for *Elthusa sinuata*. The scientific names, synonyms of parasite, and host were checked with Worms Editorial Board (2019), and Froese and Pauly (2019).



Figure 1: *Trachurus trachurus*  
(Linnaeus, 1758).



Figure 2: *Elthusa sinuata* in the gill cavity  
of *Lesueurigobius friesii* (Malm, 1874)  
(partially dissected).

## RESULTS AND DISCUSSION

Order Isopoda Latreille, 1817

Suborder Cymothoidea Wagele, 1989

Superfamily Cymothoidea Leach, 1814

Family Aegidae White, 1850

Genus *Rocinela* Leach, 1818

*Rocinela dumerilii* (Lucas, 1849) (Figs. 3-9)

Synonymies *Acherusia dumerilii* Lucas, 1849; *Acherusia complanata* Grube, 1864

*Rocinela dumerilii* Lucas, 1849: Schiodte and Meinert, 1879; Richardson, 1898; Norman, 1904; Richardson, 1905; Norman and Scott, 1906; Monod, 1923; Duncan, 1956; Menzies and Glynn, 1968; Schultz, 1969; Kensley, 1978; Kussakin, 1979; Howson and Picton, 1997; Junoy and Castelló, 2003; Alves et al., 2006; Kırkım et al., 2006; Mansour et al., 2011; Bakır et al., 2014; Bedini et al., 2015; Sampaio et al., 2016; Ortiz and Lalana, 2018.



Figure 3: *Rocinela dumerilii* (Lucas, 1849).

Host: The Atlantic horse mackerel, *Trachurus trachurus* (Linnaeus, 1758).

Infestation site: body surface. Infestation tissue: skin.

Sampling stations: 40°31'N-28°12'E; 40°33'N-27°52'E; 40°26'N-27°36'E.

**Description.** Body (14.9, 15.3, 16.9 mm) 2.4-2.6 times as long as max. width, widest at pereonite four, lateral margins subparallel. Eyes not united, separated by 9-10% width of head; each eye made up of ~13 transverse rows of ommatidia, each row with ~nine ommatidia; black eyes. Pleon with pleonite one largely concealed by pereonite seven, or visible in dorsal view; pleonite four with posterolateral margins extending beyond posterior margin of pleonite five; pleonite five with posterolateral angles acute. Pleotelson 0.8 times as long as anterior width.

Antennule peduncle article three, 0.7 times as long as combined lengths of articles one and two, 2.3 times as long as wide; flagellum with six articles, not extending to anterior of pereonite one. Antenna peduncle article three, three times as long as article two, 1.1 times as long as wide; article four 4, 1.3 times as long as article three, 1.5 times as long as wide, inferior margin with one simple seta; article five, 1.4 times as long as article four, 2.4 times as long as wide, inferior margin with one seta, anterodistal angle with cluster of three short simple setae; flagellum with 15 articles, extending to posterior of pereonite one.

Mandible molar process is a flat lobe; palp article two with seven short marginal distolateral setae and two long distolateral setae; palp article three with 16 setae. Maxillule with six RS/robust spine (one large, five slender, without serrate). Maxilla mesial lobe with two RS; lateral lobe with two RS. Maxilliped palp article one distomesial angle with three RS (with thin one seta between two large setae); article two with three hooked RS (with one thin and two large setae); article three with one hooked RS (article three fused to article two).

Pereopod one basis 2.3 times as long as greatest width; ischium three times as long as basis, inferior margin with zero RS and inferrodistal angle with two spines, superior distal margin with one seta; ischium superior distal angle with two setae and inferrodistal angle with one seta; merus inferior margin with four RS and one small spine, superior distal angle with three setae; carpus 0.7 as long as merus, inferior margin with one RS (minute); propodus 1.2 times as long as proximal width, propodal palm with blade, propodal blade 0.6 times as wide as palm, inferior margin with four RS; dactylus 1.2 times as long as propodus. Pereopods two and three similar to pereopod one (but RS on merus longer). Pereopods six and seven similar.

Pereopod seven basis 2.9 times as long as greatest width, inferior margins with 10 palmate setae; ischium 0.8 as long as basis, inferior margin with four RS, superior distal angle with three RS, inferior distal angle with four RS; merus 0.5 times as long as ischium, 2.1 times as long as wide, inferior margin with one RS, superior distal angle with five RS, inferior distal angle with four RS; carpus 0.6 as long as ischium, 2.5 times as long as wide, inferior margin with two RS, superior distal angle with eight RS (two setae), inferior distal angle with six RS; propodus 0.5 long as ischium, 4.0 long as wide, inferior margin with three RS, superior distal angle with two long and two short setae and one palme seta, inferior distal angle with two RS.

Pleopod one exopod 1.9 times as long as wide, with PMS/plumose seta on right lateral margin, with PMS on two-thirds of left lateral margin; mesial margin weakly convex, with PMS; endopod 2.2 times as long as wide, lateral margin weakly concave, with PMS on two-third of left margin, without PMS on right margin; mesial margin with PMS; peduncle mesial margin with five-six coupling hooks. Pleopods two-four peduncle mesial margin with five-six coupling hooks, lateral margin with PMS. Uropod peduncle ventrolateral margin with two RS and five-seven PMS, posterior lobe half long as endopod. Exopod rami not extending to pleotelson apex. Endopod lateral margin convex, lateral margin with five RS, mesial margin distally rounded, with five RS. Exopod not extending to endopod end, 3.2 times long as maximum width; lateral margin convex, with eight RS; mesial margin straight, distally convex, with zero RS; distal margin with indistinct apex.




Figure 4: *Rocinela dumerilii* a. antenna (1.31 mm), b. antennule (0.8 mm), c. maxilla (0.16 mm), d. maxillule (0.37 mm), e. maxilliped (0.37 mm), f. mandible (0.57 mm).




Figure 5: *Rocinela dumerilii* (a-g),  
Pereopods 1-7 (0.66 mm).






Figure 6: *Rocinela dumerilii* (a-e), Pleopods 1-5 (1.21 mm).




Figure 7: *Rocinela dumerilii*, Uropod and apex of exopod.




Figure 8: *Rocinela dumerilii*, head.



Figure 9: *Rocinela dumerilii*, penes.

**Remarks.** Forty-one species belonging to *Rocinela* genus are listed by Worms Editorial Board (2019). The monographs, detailed descriptions, and revision studies concerning *Rocinela* genus have been published mostly on the Australia, New Zealand, and Pacific Ocean (Bruce, 1983; 2009; Brusca, 1983; Brusca and France, 1992).

The historic literature about the species of *Rocinela* in the Mediterranean Sea lack modern or detailed descriptions (Schiodte and Meinert, 1879; Kussakin, 1979; Sars, 1899). Three species belonging to *Rocinela* genus were reported such as *Rocinela ophthalmica* Milne Edwards, 1840, *Rocinela dumerilii* (Lucas, 1849), and *Rocinela danmoniensis* Leach, 1818 from the Mediterranean Sea.

Although there are no detailed descriptions currently available to compare the morphological characters of *R. dumerilii* and *R. danmoniensis*, some morphological characters such as the setation on propodus of pereopods and the distance between eyes were compared in this study. *R. dumerilii* species is very closely similar to *R. Danmoniensis* species. The main point of difference is that in *R. danmoniensis* the eyes are closer together and may even touch other (Schiodte and Meinert, 1879; Norman, 1904; Richardson, 1905; Norman and Scott, 1906; Schultz, 1969; Kussakin, 1979; Kırkım et al., 2006). The eyes on *Rocinela* samples in this study are separated by about 9-10% width of head. The propodus of the pereopod two bears four setae in female and male of *R. dumerilii*, but two setae in the propodus of the pereopod one of pullus primus of *R. danmoniensis* according to Schiodte and Meinert (1879). The setation of propodus in this study is consistent with *R. dumerilii*. The morphological extremities of mouth-parts of specimens found in this study have been compared to the findings presented by Monod (1923). The morphology of specimens collected in this study is consistent with the literature concerning *R. dumerilii* (Schiodte and Meinert 1879; Monod, 1923; Kussakin, 1979).

There are several association records between isopods belonging *Rocinela* genus and fish around the world (Novotny and Mahnken, 1971; Brusca and Iverson, 1985; Kensley and Schotte, 1989; Garzón-Ferreira, 1990; Williams et al., 1994; Wing and Moles, 1995; Bunkley-Williams and Williams, 1998; De Lima et al., 2005). Regarding the Mediterranean Sea, there were micropredator reports of *R. danmoniensis* from fish as host. *R. danmoniensis* was reported on Atlantic cod, *Gadus morhua* (Gadidae) (Schiodte and Meinert, 1879) and Blackspot seabream, *Pagellus bogaraveo* (Sparidae) (Hermida et al., 2013). Sars (1899) mentioned that it is more generally found clinging to the skin of fish of various kinds, for instance the common cod, the haddock, the ling, etc. Stephensen (1948) indicated the occurrence of its parasitism on Gadidae such as cod, haddock and lange.

There were some reports of *Rocinela dumerilii* surviving as free-living organisms and from the digested stomach content of some marine organisms. Bedini et al. (2015) reported this species among macroinvertebrate assemblages within a *Posidonia oceanica* meadow. Sampaio et al. (2016) recorded it at 48 meters depth of the Portuguese continental shelf. Alves et al. (2006) recorded it in stomach contents of *Sepia officinalis* Linnaeus, 1758 (Mollusca, Cephalopoda) off the south coast of Portugal, while this species was also reported among the food of grey seals, *Halichoerus grypus* (Fabricius, 1791) (Chordata, Mammalia) from the Isle of Man by Duncan (1956), and as food of the Almaco Jack, *Seriola rivoliana* Valenciennes, 1833 (Chordata, Actinopterygii) in Tunisian marine waters by Mansour et al. (2011).

*R. dumerilii* was collected on the *Trachurus trachurus* body in this study. *Trachurus trachurus* (Linnaeus, 1758) (Perciformes; Carangidae) is pelagic-neritic and oceanodromus fish with commercial value. It is one of the most important pelagic fish of purse seine and trawling fishery (Froese and Pauly, 2019). The total amount of european hake has been decreasing about 22.200 tones in 2000 to 12.985 tones for 2017 in Turkey (TUIK, 2018).

There is no report regarding micropredations of *R. dumerilii* on fish or other marine organisms according to the available literature. When the interrelationships between members of Carangidae as host and Aegidae as micropredator were examined, the occurrence of some records in the literature is supportive of our finding micropredator behaviour of *R. dumerilii* being also aegid on the atlantic horse mackerel in this study. Bruce (2009) found *Aegapheles alazon* species from the yellowtail horse mackerel, *Trachurus novaezelandiae* Richardson, 1843, belonging to Carangidae family in New Zealand. *R. signata* was reported on *Caranx crysos* (Mitchill, 1815) (Carangidae) and *Oligoplites saliens* (Bloch, 1793) (Carangidae) from Bahia by Carvalho-Souza et al. (2009), while it was reported on *Chloroscombrus chrysurus* (Linnaeus, 1766) (Carangidae) from Rio Grande do Norte by Lima et al. (2011); on *Caranx* sp. from Jamaica; Carrie Bow Cay and Blue Ground Range, in Belize by Kensley and Schotte (1989).

Mansour et al. (2011) found that *Rocinela dumerilii* from the stomach content of almaco jack, *Seriola rivoliana*, being carangid fish in Tunisian marine waters. Torcu Koç and Erdoğan (2019) and Bayhan et al. (2013) indicated that crustaceans were the dominant food category of atlantic horse mackerel. It is our opinion that Atlantic horse mackerel, being a migratory and carnivorous fish species, may have interacted with these isopods during their feeding activities during migrations.

Stepien and Brusca (1985) reported that fish were attacked by members of demersal crustacean assemblage such as luminent ostracod, *Vargula tsujii*, and the isopod *Cirolana diminuta* during their night diving observations while studying the case of death of adult fish in large cages on the seafloor off of the Southern California coast. They summarized the relationships and interactions between nocturnal migrating zooplankton and fish throughout the world.

Nocturnal migrating zooplankton assemblages consist of crustaceans including copepods, ostracods, amphipods, isopods, mysids, cumaceans, decapod larvae, and shrimps. These zooplankton assemblages pose a predation risk for migrating fish in comparison with non-migrating fish. Most nocturnally-active reef fish feed primarily on these migrating forms (Stepien and Brusca, 1985).

Stepien and Brusca (1985) indicated that ostracods are more attracted to chemicals released by healthy, sexually mature fish in comparison to juvenile fish. The chemicals from fish injured by ostracods may serve to attract amphipods and isopods. They observed that both ostracods and isopods often invaded the gills, entered the body cavity, and consumed the fish' gonads and liver. Stepien and Brusca (1985) also discussed that whether anecdotally reported these attacks occurred on live, dying or already-dead fish in the Pacific Ocean, off the coasts of Japan and the Arctic. This may be a possible reason, needed to be studied, for these attacks as the chemicals released from wounds being on fish after caught in the gill net attract isopods and amphipods. It is implied that the phenomenon about attack behaviour may be common in most coastal areas in the world. This attack behaviour was also observed by Turkish commercial fishermen many times during our research surveys.

Family Cymothoidae Leach, 1814

*Elthusa* Schioedte and Meinert, 1884

*Elthusa sinuata* (Koelbel, 1879) (Fig. 8)

Synonyms: *Livoneca mediterranea* Heller, 1868; *Livoneca sinuata* Koelbel, 1879; *Livoneca sinuata* Brian, 1912.



Figure 8: *Elthusa sinuata* (1 mm).

Host: the Fries's goby, *Lesueurigobius friesii*.

Infestation area, gill cavity. Examined fish, 70. Infested fish, five. Prevalence, 7.12%. Mean intensity, one.

Sampling stations: 40°31'N-28°12'E; 40°33'N-27°52'E; 40°33'N-27°41'E; 40°26'N-27°36'E.

*Elthusia sinuata* (syn. *Livoneca sinuata*) was described on the red bandfish, *Cepola macrophthalma* (syn. *Cepolae rubescentis*) from Sicilia by Koelbel (1879). It was reported from North-West Africa, United Kingdom, Mediterranean and Adriatic seas, Spain, France, Algeria, Tunisia, Italy, Yugoslavia, Montenegro, Turkey (van der Wal et al., 2019).

This parasitic isopod has been collected on hosts from several fish species as *Cepola macrophthalma* (Cepolidae), *Pleuronectes* sp. (Pleuronectidae), *Raja miraletus* (Rajidae), *Boops boops* (Sparidae), *Gobius* sp. (Gobiidae), *Brama brama* (Bramidae), *Lepidopus caudatus* (Trichiuridae), *Trichiurus lepturus* (Trichiuridae), *Raja clavata* (Rajidae), and *Argentina sphyraena* (Argentinidae). It was also recorded from two cephalopod species *Sepiolo ligulata* and *Loligo vulgaris* except of fish (Öktener et al., 2018; van der Wal et al., 2019).

Although no comment can be made about the host selectivity of *Elthusia sinuata*, six of the ten parasitized species belong to Perciformes. Most of host species exhibit demersal character and carnivorous feeding behaviour. This parasite is found on the gill cavity of Fries's goby, *Lesueurigobius friesii* in this study. This new host species also exhibits demersal character and carnivore feeding behaviour. The characters of this gobiid are similar to the characters of other hosts. This is consistent with findings in *Gobius* sp. reported by Dollfus and Trilles (1976).

### CONCLUSIONS

Members of Aegidae are known for their behavior as micropredators or temporary parasites. Although species of these micropredators from Mediterranean have been reported by several researchers many times, information about the biology, life cycles, and interactions and relationships with other organisms of these micropredators are limited. The aim of this study is to contribute scientific information regarding the aforementioned unknown areas about these micropredators.

### ACKNOWLEDGEMENTS

The samples identified in this study were obtained in the sampling area of study for (TAGEM/HAYSUD/2014/05/01) Project carried by Sheep Research Institute and supported by Ministry of Food, Agriculture and Livestock of Turkey.

## REFERENCES

1. Alves D. M., Cristo M., Sendao J. and Borges T. C., 2006 – Diet of the cuttlefish *Sepia officinalis* (Cephalopoda: Sepiidae) off the south coast of Portugal (eastern Algarve), *Journal of the Marine Biological Association of the United Kingdom*, 86, 2, 429-436, doi: 10.1017/S0025315406013312.
2. Bakır A. K., Kayağan T., Aker H. V., Özcan T., Sezgin M., Ateş A. S., Koçak C. and Kırkım F., 2014 – The marine arthropods of Turkey, *Turkish Journal of Zoology*, 38, 765-831, doi:10.3906/zoo-1405-48.
3. Bayhan B., Sever T. M. and Kara A., 2013 – Diet composition of the Mediterranean horse mackerel, *Trachurus mediterraneus* (Steindachner, 1868) (Osteichthyes: Carangidae), from the Aegean Sea, *Belgian Journal of Zoology*, 143, 15-22.
4. Bedini R., Bedini M., Bonechi L. and Piazzini L., 2015 – Patterns of spatial variability of mobile macro-invertebrate assemblages within a *Posidonia oceanica* meadow, *Journal of Natural History*, 49, 2559-2581, doi: 10.1080/00222933.2015.1021872.
5. Bruce N. L., 1983 – Aegidae (Isopoda: Crustacea) from Australia with descriptions of three new species, *Journal of Natural History*, 17, 757-788, doi: 10.1080/00222938300770591.
6. Bruce N. L., 2003 – Giant marine bloodsuckers, *Water and Atmosphere*, 11, 3, 4.
7. Bruce N. L., 2004 – Reassessment of the isopod crustacean *Aega deshaynsiana* (Milne Edwards, 1840) (Cymothoidea: Aegidae): a worldwide complex of 21 species, *Zoological Journal of the Linnean Society*, 142, 2, 135-232, doi: 10.1111/j.1096-3642.2004.00127.x.
8. Bruce N. L., 2009 – The Marine Fauna of New Zealand: Isopoda, Aegidae (Crustacea), *NIWA Biodiversity Memoir*, 122, 1-252.
9. Brusca R. C., 1983 – A monograph on the isopod family Aegidae in the tropical eastern Pacific, The genus *Aega*, *Allan Hancock Foundation Monographs in Marine Biology*, 12, 1-39.
10. Brusca R. C. and Iverson E. W. 1985 – A Guide to the marine Isopod Crustacea of Pacific Costa Rica, *Revista de Biología Tropical*, 33, 1-67.
11. Brusca R. C. and France S. C., 1992 – The genus *Rocinela* (Crustacea: Isopoda: Aegidae) in the tropical eastern Pacific, *Zoological Journal of the Linnean Society*, 106, 231-275, doi: 10.1111/j.1096-3642.1992.tb01248.x.
12. Bunkley-Williams L. and Williams J. R. E. H., 1998 – Isopods associated with fish: a synopsis and corrections, *Journal of Parasitology*, 84, 5, 893-896, doi: 10.2307/3284615.
13. Carvalho-Souza G. F., Souza Neto J. R., Aleluia F. T., Nascimento I. A., Browne-Ribeiro H., Santos R. C. and Tinoco M. S., 2009 – Occurrence of isopods ectoparasites in marine fish on the Cotegipe Bay, north-eastern Brazil, *Marine Biodiversity Records*, 2, 1-4, doi: 10.1017/S1755267209990844.
14. Castelló J. and Junoy J., 2003 – Catálogo de las especies ibéricas y baleares de isópodos marinos (Crustacea: Isopoda), *Boletín del Instituto Español de Oceanografía*, 19, 293-326. (in Spanish)
15. De Lima J., Chellappa S. and Thatcher V. E., 2005 – *Livoneca redmanni* Leach (Isopoda, Cymothoidea) and *Rocinela signata* Schioedte and Meinert (Isopoda, Aegidae), ectoparasites of *Scomberomorus brasiliensis* Collette, Russo and Zavala-Camin (Osteichthyes, Scombridae) no Rio Grande do Norte, Brazil, *Revista Brasileira de Zoologia*, 22, 4, 1104-1108, doi: 10.1590/S0101-81752005000400041.
16. Dollfus R. P. and Trilles J. P., 1976 – A propos de la collection R. P. Dollfus, mise au point sur les cymothoïdiens jusqu'à présent récoltés sur des téléostéens du Maroc et de l'Algérie, *Bulletin du Muséum d'histoire naturelle Paris*, 272, 821-830. (in French)
17. Duncan A., 1956 – Notes on the food and parasites of two Grey Seals, *Halichoerus grypus* (Fabricius) from the Isle of Man, *Journal of Zoology*, 126, 4, 635-644, doi: 10.1111/j.1096-3642.1956.tb00455.x.
18. Froese R. and Pauly D. (eds), 2019 – FishBase, World Wide Web electronic publication, www.fishbase.org, version (02/2019).
19. Garzón-Ferreira J., 1990 – An isopod, *Rocinela signata* (Crustacea: Isopoda: Aegidae), that attacks humans, *Bulletin of Marine Science*, 46, 813-815.

20. Hermida M., Cruz C. and Saraiva A., 2013 – Parasites as biological tags for stock identification of blackspot seabream, *Pagellus bogaraveo*, in Portuguese north-east Atlantic waters, *Scientia Marina*, 77, 4, 607-615, doi: 10.3989/scimar.03859.17A.
21. Howson C. M. and Picton B. E., 1997 – The species directory of the marine fauna and flora of the British Isles and surrounding seas, Ulster Museum Publication, 276, The Ulster Museum: Belfast, U.K., ISBN 0-948150-06-8. vi, 508.
22. Kensley B., 1978 – Guide to the Marine Isopods of Southern Africa, South African Museum and The Rustica Press, Wynberg, Cape Town, 173.
23. Kensley B. and Schotte M., 1989 – Guide to the marine isopod crustaceans of the Caribbean, Smithsonian Institution Press, Washington D.C., 308.
24. Kırkım F., Kocataş A., Katağan T. and Sezgin M., 2006 – Contribution to the knowledge of the free-living Isopods of the Aegean Sea Coast of Turkey, *Turkish Journal of Zoology*, 30, 361-372.
25. Koelbel C., 1879 – Über einige neue Cymothoiden, *Sitzungsber Kaiserl Akad Wiss, Math-Naturwiss*, 1, 78, 401-416. (in German)
26. Kussakin O. G., 1979 – Marine and brackishwater likefooted Crustacea (Isopoda) from the cold and temperate waters of the Northern Hemisphere, Suborder Flabellifera, Izdatel'stvo Nauka, Leningrad, *Opredeliteli po Faune S.S.S.R., Izdavaemye Zoologicheskim Institutom Akademii Nauk S.S.S.R.*, 472. (in Russian)
27. Lima J., Freitas M. D. F., Fernandes B. L. F. and Bezerra J. T. A., 2011 – Preferência de diferentes habitats do parasita *Rocinela signata* em peixes marinhos *Archosargus rhomboidalis* e *Chloroscombrus chrysurus* no litoral do Rio Grande do Norte, *Revista Centauro*, 2, 23-27. (in Portuguese)
28. Mansour S., Azzouz K., Boumaïza M., Ben Amor M. M. and Capapé C., 2011 – First record of a rare carangid species, the Almaco Jack, *Seriola rivoliana* (Osteichthyes: Carangidae) in Tunisian marine waters (central Mediterranean), *Cahiers de Biologie Marine*, 52, 187, 192.
29. Menzies R. and Glynn P. W., 1968 – The common marine isopod Crustacea of Puerto Rico, *Studies on the Fauna of Curaçao and other Caribbean Islands*, 104, 1-133.
30. Monod T., 1923 – Prodrome d'une faune des Tanaidacea et des Isopoda (excl. Epicaridea) des côtes de France (excl. Méditerranée), *Annales de la Société des Sciences Naturelles de la Charente-Inférieure*, 33, 19-125. (in French)
31. Howson C. M. and Picton B. E., 1997 – The species directory of the marine fauna and flora of the British Isles and surrounding seas, Ulster Museum Publication, 276, The Ulster Museum: Belfast, U.K., ISBN 0-948150-06-8. vi, 508.
32. Norman A. M., 1904 – British Isopoda of the families Aegidae, Cirolanidae, Idoteidae and Arcturidae, *Annals and Magazine of Natural History*, 7, 430-450, doi: 10.5962/bhl.title.35831.
33. Norman A. M. and Scott T., 1906 – The Crustacea of Devon and Cornwall, London: William Wesley and Son. 232, doi: 10.5962/bhl.title.1939.
34. Novotny A. J. and Mahnken C. V. W., 1971 – Predation on juvenile Pacific salmon by *Rocinela belliceptis pugettensis* (Crustacea: Isopoda), *Fishery Bulletin*, 69, 699-701.
35. Ortiz M. and Lalana R., 2018 – Lista de especies y distribución de los isópodos (Crustacea: Peracarida) de Cuba, *Novitates Caribaea*, 12, 102-126, doi: 10.33800/nc.v0i12.87. (in Spanish)
36. Öktener A., Trilles J. P., Kocabaş E., İnce H. and Alaş A., 2018 – Redescription of *Elthusa sinuata* (Koelbel, 1879) comb. nov. (Isopoda, Cymothoidae) parasitizing the red bandfish in Turkey, *Thalassas*, 34, 333-339, doi: 10.1007/s41208-018-0068-z.
37. Ramdane Z. and Trilles J. P., 2008 – Cymothoidae and Aegidae (Crustacea, Isopoda) from Algeria, *Acta Parasitologica*, 53, 173-178, doi: 10.2478/s11686-008-0033-8.
38. Richardson H., 1898 – Description of four new species of "Rocinela", with a synopsis of the Genus, *Proceedings of the American Philosophical Society*, 37, 157, 8-17, doi: 10.2307/983687.
39. Richardson H., 1905 – A monograph on the isopods of North America, *Bulletin of United States National Museum*, 54, 1-727. doi: 10.5479/si.03629236.54.



40. Sampaio L., Mamede R., Ricardo F., Magalhaes L., Rocha H., Martins R., Dauvin J. C., Rodrigues A. M. and Quintino V., 2016 – Soft-sediment crustacean diversity and distribution along the Portuguese continental shelf, *Journal of Marine Systems*, 163, 43-60, doi: 10.1016/j.jmarsys.2016.06.011.
41. Sars G. O., 1899 – An account of the Crustacea of Norway with short descriptions and figures of all the species, II, Isopoda, 13, 14, Bergen Museum, Christiania, 264, doi: 10.5962/bhl.title.1164.
42. Schiodte J. C. and Meinert F., 1879 – Symbolae ad monographiam Cymothoarum, Crustaceorum, Isopodum Familiæ, I, Aegidae, *Naturhistorisk Tidsskrift, Tredie Raekke*, 12, 321-414, doi: 10.5962/bhl.title.10300.
43. Schultz G. A., 1969 – The marine isopod crustaceans, How to know, The pictured key, Nature series, WMC Brown Company Publishers, Dubuque, Iowa, 1-359.
44. Stephensen K., 1948 – Storkrebs IV, 3, Tanglus (Marine Isopoder) og Tanaider, Danmarks Fauna 53, København: G. E. C., Gads Forlag, 187. (in Danish)
45. Stepien C. A. and Brusca R. C., 1985 – Nocturnal attacks on nearshore fish in southern California by crustacean zooplankton, *Marine Ecology Progress Series*, 25, 91-105, doi: 10.3354/meps025091.
46. Torcu Koç H. and Erdoğan Z., 2019 – Some population parameters of the Mediterranean horse mackerel, [(*Trachurus mediterraneus* (Steindachner, 1868)] population living in Bandırma Bay, The Sea of Marmara, *Journal of BAUN Institute Science Technology*, 21, 1, 265-277.
47. Treasurer J. W., 2001 – A parasitic marine isopod *Aega tridens* Leech on mackerel *Scomber scomber* L. in British Waters, *Bulletin of the European Association of Fish Pathologists*, 21, 1, 37-39.
48. Trilles J. P., 1976 – Les Cymothoidae (Isopoda, Flabellifera) des collections du Museum National d'Histoire Naturelle de Paris, IV, Les Lironecinae Schioedte and Meinert, 1884, *Bulletin du Muséum d'histoire naturelle Paris*, 272, 773-800. (in French)
49. TUIK, 2018 – Fishery Statistics, available from <http://www.tuik.gov.tr/Start.do>.
50. van der Wal S., Smit N. J. and Hadfield K A., 2019 – Review of the fish parasitic genus *Elthusa* Schioedte and Meinert, 1884 (Crustacea, Isopoda, Cymothoidae) from South Africa, including the description of three new species, *Zookeys*, 23, 1-37, doi: 10.3897/zookeys.841.32364.
51. Williams J. R. E. H., Bunkley-Williams L. and Sanner C. J., 1994 – New host and locality records for copepod and isopod parasites of Colombian marine fish, *Journal of Aquatic Animal Health*, 6, 362-364.
52. Wing B. A. and Moles D. L., 1995 – Behavior of *Rocinela angustata* (Isopoda, Aegidae), an ectoparasite of Alaskan marine fish, *Journal of Aquatic Animal Health*, 7, 1, 34-37.
53. WoRMS Editorial Board 2019 – World Register of Marine Species, available from <http://www.marinespecies.org> at VLIZ, accessed in 2019-07-23, doi:10.14284/170.