

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/344197617>

Acute toxicity of two pesticides (dursban, mancozeb) and their combined mixture on the terrestrial isopod *Armadillidium vulgare* (Oniscidea, Isopoda)

Article · January 2019

CITATION

1

READS

28

11 authors, including:



Maya Moumene

Université de Béjaïa

2 PUBLICATIONS 1 CITATION

[SEE PROFILE](#)



Mohamed Nait Mouloud

Université Mouloud Mammeri de Tizi Ouzou

9 PUBLICATIONS 19 CITATIONS

[SEE PROFILE](#)



Mahdeb Amokrane

Inraa

10 PUBLICATIONS 6 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Valorization of Prickly Pear (*Opuntia ficus indica*) products and by-product [View project](#)



Manca stages of terrestrial isopods (Crustacea, Isopoda, Oniscidea) [View project](#)

Acute toxicity of two pesticides (dursban, mancozeb) and their combined mixture on the terrestrial isopod *Armadillidium vulgare* (Oniscidea, Isopoda)

Karima Hachemaoui Benmouhoub^{1*}, Maya Moumene¹, Chafika Mouhoub Sayah², Mohamed Nait Mouloud³, Hafsa Djoudad Kadji¹, Saddek Yesguer¹, Mokrane Mahdeb⁴, Aouba Mouzaia Salhi¹, Faouzia Charfi⁵, Giuseppe Montesanto⁶, Caroline Habold⁷

¹Laboratory of Applied Zoology and Animal Ecophysiology, Faculty of Natural and Life Sciences, University of Bejaia, 06000 Bejaia, Algeria

²University of Bouira, Algeria

³Laboratory in Marine and Aquaculture Ecosystems, Faculty of Natural and Life Sciences, University of Bejaia, 06000, Bejaia, Algeria

⁴Mountain Systems Agro Division, INRA, Oued Ghir Bejaia, Algeria

⁵Laboratory of Biology and Evolutionary Systems, Faculty of Sciences, El Manar University, Tunis.

⁶ Department of Biology, Pisa, Italy

⁷Institut Pluridisciplinaire Hubert CUREN STRASBOURG, France

Abstract: Our study focused on evaluating an acute toxicity test, where woodlice (*Armadillidium vulgare*) which is the most abandoned in our region, were exposed to pesticides (dursban and mancozeb) most used in Algeria. The concentrations of the pesticides tested are 720 mg/kg, 360 mg/kg, 90 mg/kg, 45 mg/kg for dursban; 800 mg/kg, 400 mg/kg, 200 mg/kg, 100 mg/kg for mancozeb and 720/800 mg/kg, 360/400 mg/kg, 90/200 mg/kg, 45/100 mg/kg for the mixture of dursban and mancozeb. Our results show that dursban was highly toxic (69% mortality after 28 days of the test) compared to mancozeb (19.5% mortality). On the other hand, in the case of the mixture (dursban ± mancozeb), the toxicity was lower (63% mortality) compared to the dursban tested alone. Contamination of woodlice with dursban and a mixture of dursban and mancozeb revealed morphological effects such as alteration of the marsupial cavity and expulsion of eggs from the outside of this cavity. Dursban and the mixture of dursban and mancozeb are toxic to *Armadillidium vulgare*, but mancozeb alone is not toxic. Dursban should be banned because it is bad for soil fauna and therefore probably for biodiversity in general and even human health.

Keywords: bio indicator, dursban, mancozeb, acute toxicity, *Armadillidium vulgare*.

INTRODUCTION

Contamination and then pollution of the soil, by various factors, directly causes serious consequences on living beings and indirectly on the environment. The development of organic chemistry marks the arrival (in 1960) of synthetic "pesticide" molecules that are widely used to improve agricultural production due to food needs that are constantly increasing with population growth (Pretty, 2008). As a result, Algeria imported an average of 17159,648 tonnes of pesticides during 2017 at an estimated cost of US\$ 855,38253.82 (ASISD, 2017).

The Soummam Valley, which is an agricultural area of 130,348 hectares, is known for its significant agricultural activity, particularly market gardening and arboriculture. A total of 1945,077 tonnes of pesticides during 2017 was used for these activities (ASB, 2017). The survey of farmers in this region showed that mancozeb and dursban are the most commonly used pesticides in agricultural practices due to their availability on the market and their proven efficiency against pests. However, the systematic use of these products is currently being questioned, with growing awareness of the risks they can generate. One of the most effective approaches for assessing the effects of pesticides in the environment and therefore the risk associated with their contamination is bio-monitoring (Salines, 2012).

This bio monitoring principle has been used on various bio-indicators of soil pollution such as earthworms (Annelid) (Vermeulen et al., 2001), rats (Szepvolgyi et al., 1989), snails (Coline, 2011; Viard, 2004), and aquatic crustaceans (Palma et al., 2008). We add to this list of bio-indicators of soil quality, some Oniscidea species such as terrestrial isopods or woodlice (Grosset et al., 2005; Kammenga et al., 2000) which are abundant in various terrestrial ecosystems (Cortet et al., 1999; Camila et al., 2017) in which they constitute an essential link in the food web (Bureš et al., 2003). They contribute enormously in increasing soil fertility (Hassall et al., 1978; Curry, 1994; Camila, 2014). As such they can be used as bio-indicators for the eco-toxicological analysis of pesticide pollution (Van Gestel et al., 2018). It is in this context that the present work aims to evaluate the effect of the two most widely used pesticides i.e. mancozeb and dursban in the Soummam Valley in Algeria, on the survival and morphology of males and females *Armadillidium vulgare*, the most abundant species in the study site.

Several studies have highlighted the effects of the two pesticides on the mortality of terrestrial isopod species (Morgado et al., 2016; Morgado et al., 2018; Jänschet al., 2005), but no acute pesticide toxicity studies have been carried out yet on the species *Armadillidium vulgare*. Also, this is the first study

investigating the effects of these pesticides on the morphology of these species.

MATERIALS AND METHODS

Study station

The Soummam Valley region extends for about 120 km from the southwestern slope of the Djurdjura mountain range (latitude 36°21'8 N; longitude 3°53'E) to the coast of the Mediterranean Sea. It is a socio-economic regional entity in the North East of Algeria. It is an agricultural area where the cultivated plots are located along side urban centers and scattered hamlets. It is covered by a fairly dense hydrographic network. It is also characterized by atypically. Mediterranean vegetation, composed mainly of bulges, cultivated plants and maquis of arborescent strata (cork oak, holm oak, Aleppo pine, poplar and olive tree). The climate is Mediterranean with hot, dry summers and cold, rainy winters.

Biological model

In this experiment, the terrestrial isopod *Armadillidium vulgare* was used as a test species. These organisms were collected by the National Institute of Agricultural Research, Algeria (INRA) and then maintained in the laboratory of Applied Zoology and Animal Ecophysiology at room temperature (22°C±1°C), and a photoperiod of 13h/11h (light/dark). Only adult isopods (7 pairs of periopods), a body mass of 15-25 mg were used in this experiment. The individuals selected in this test were healthy.

Chemical compound and soils

Two pesticides were used for this experiment which commercial formulations are: 1/ the organophosphorus insecticide dursban that contains 480g/l of chlorpyrifos (DOW AGRO SCIENCE) and is used to treat many pests such as caterpillars and sucking insects (mealy bugs, aphids) and 2/ the dithiocarbamate fungicide, mancozeb with 80% of mancozeb (SHANGHA, DEZENTELU International TRADE CO); used for fruit tree treatments for example against apple scab.

The soil used for this biotest was taken from the *Armadillidium vulgare* sampling station and then subjected to physico-chemical soil analyses. The main properties of this soil are: pH = 8.56 ± 0.05 (pHmeter), total limestone = 3.6 ± 0.2 (%) (Bernard's Calcimeter); texture (Robinson's pipette grain size): 24 ± 1.08 (%) clay; 24 ± 1.53 (%) silt; 52 ± 2 (%) sand. Thus the soil is sandy-clay in nature according to the Atterberg triangle textures. The soil was heated to 80°C for several hours to destroy predators before isopods were introduced.

The experimental protocol

In these bioassays, the concentrations of both pesticides (dursban and mancozeb) were used in mg of active ingredient per kg of soil as used previously on

earthworms (OECD, 1984; ISO, 2008). For this acute toxicity test, we took four different concentrations for each of the pesticides (dursban and mancozeb). The highest concentrations were those used by the farmer in the field (720 mg/kg for dursban and 800 mg/kg for mancozeb). The three other concentrations were below the maximum concentrations (720 mg/kg, 800 mg/kg) in decreasing logarithmic order. The doses of pesticides used were: & dursban: 0 mg/kg (control), 720 mg/kg, 360 mg/kg, 90 mg/kg, 45 mg/kg. & mancozeb: 0 mg/kg (control), 800 mg/kg, 400 mg/kg, 200 mg/kg, 100 mg/kg. & Binary mixture (dursban ± mancozeb): 0/0 mg/kg (control), 720/800 mg/kg, 360/400 mg/kg, 90/200 mg/kg, 45/100 mg/kg. The test was repeated 4 times for each of these concentrations. Thus the experimental protocol adopted for our bioassay required the preparation of 60 plastic boxes of the same dimensions (30cm x 20cm x 10cm), with perforated lids for ventilation. Contamination of the environment was carried out by spraying dilutions of prepared pesticides over the entire surface of the soil placed in the boxes. After impregnation with the pesticide for 2 hours, ten adults (7 pairs of periopods) woodlice of each sex (5 males and 5 females) were introduced into each box. Woodlice were fed daily on litter collected from the sampling site composed of cork oaks, holm oaks, Aleppo pines, dry poplars, olive and carrot, potato peels and water. Mortality control was performed once a week. Dead woodlice were counted after 7, 14, 21 and 28 days.

Morphological effects of pesticides on *Armadillidium Vulgare*

During this experiment we looked for other morphological effects caused by the pesticides tested, by identifying all the morphological anomalies observed on dead or alive woodlice.

Statistical analysis

The statistical analysis was performed with R software package version 3.5.1 (2018) in GLMS. A post Hoc test (Tuckey method) was performed to compare between doses using the emmeans function in emmeans library.

Lethal concentration was determined using R software by Koissi Saviet al (2017).

RESULTS

Acute toxicity test on the species

Armadillidium vulgare

Effect of dursban on the survival of terrestrial isopods *Armadillidium Vulgare*

After 28 days, the effect of dursban doses on *Armadillidium Vulgare* was significant ($p < 0.05$). A proportional relationship was observed between the doses used and the mortality obtained. (Fig.1). After already 7 days of exposure, the highest doses of dursban (720 mg/kg) caused 100% mortality in the population. The same result was observed after 14 days at a concentrations of 360 mg/kg.




Fig. 1 Survival at 28 days (mean \pm sd) of *Armadillidium vulgare* after exposure to single treatment of dursban. Different letters (a,b,c,d) indicate significant differences between concentrations ($P < 0.05$).

Effect of mancozeb on the survival of terrestrial isopods *Armadillidium Vulgare*

Mancozeb induced a lower mortality in woodlice, compared to dursban. However a significant decrease in survival ($p < 0.05$) was observed at 28 days at the concentrations of 200-800 mg/kg compared to the control condition. (Fig.2).

During the first week, the mortality rate of woodlice subjected to mancozeb was 5.12% for all concentrations (800,400, 200,100 mg/kg). At 28 days, the mortality was 18.5 % at the highest concentration (800 mg/kg) and 12.5 % at the lowest concentration (100 mg/kg).




Fig. 2 Survival (mean \pm sd) of *Armadillidium vulgare* after a 28 days exposure to single treatment of mancozeb. Different letters (a,b) indicate significant differences between concentrations ($P < 0.05$).

Effect of the mixture (dursban \pm mancozeb) on the survival of terrestrial isopods *Armadillidium Vulgare*

The effect of all doses was significant compared to the control. The concentration 45/100 and 90/200 mg/kg had the same effect ($p < 0.05$) but beyond these doses, mortality became more and more significant. The

dose 720/800 mg/kg caused the highest mortality. (Fig.3.) After already one week of exposure (7 days), the highest doses of the mixture (720/800 mg/kg) caused 100% of mortality in the population. The same result was observed for the concentrations 360/400 mg/kg after 14 days. After 28 days of exposure the lowest dose (45/100 mg/kg) revealed 30% of mortality.




Fig. 3. Survival (mean \pm sd) of *Armadillidium vulgare* after a 28 day exposure to a mixture of mancozeb and dursban. Different letters (a, b, c,d) indicate significant differences between concentrations ($P < 0.05$).

Determination of the LD50

The LD50 (the lethal dose of 50% of the population) of dursban after 14 days of experiment was 72.77 mg/kg. (Fig.4). Determination of the LD50 was not possible in an acute mancozeb toxicity test because

mortality was not greater than or equal to 50% (25% mortality in the experimental population). The LD50 of the binary mixture after 14 days of experiment was 330 mg/kg (Fig.5).




Fig. 4 Dose response curves for the acute toxicity test with *Armadillidium vulgare* in dursban (LD50).




Fig. 5 Dose response curves for the acute toxicity test with *Armadillidium vulgare* submitted to a mixture of dursban \pm mancozeb (LD50).

Morphological effects of pesticides on *Armadillidium Vulgare*

In the acute toxicity test conducted on woodlice, some morphological anomalies were observed in the individuals tested with dursban and the mixture of

dursban ± mancozeb: *Paralysis of the anterior part in some individuals (Fig.6a). *Marsupium destruction and mucus release (Fig.6b). *Egg expulsion outside the marsupium (Fig.6c).



Fig. 6. Anomalies observed in woodlice contaminated by dursban, and the mixture dursban ± mancozeb: (a) paralysis of anterior part, (b) destruction of marsupium and release of mucus, (c) expulsion of eggs from marsupium.

DISCUSSION

After one month of acute toxicity testing on the *Armadillidium vulgare* population, a low toxicity of mancozeb was recorded (mortality of 19.5%). The same concentrations of this pesticide were tested by Yesguer (2015) on *Aporrectodea caliginosa* (Annelid species). These results corroborate those of our study. Also, Vermeulen et al (2001) and Ioriatti et al (1992) confirmed this low mancozeb toxicity respectively in the species *Eisenia Andrei* (Annelid) and the predatory mite *Amblyseius andersoni*.

Mancozeb had even no effect on adult *Amblyseius fallacies* (Acarid species) (Bostanian et al.,1998). These results are explained by the low solubility of this fungicide in lipids. According to Fabre et al (1954) this product accumulates poorly in animal tissues and therefore is easily diluted in water and dispersed after rain or irrigation. The LD50 for mancozeb was not determined because mortality was less than 50% (19.5%). Similar values were obtained by Yesguer 2015 on *Aporrectodea caliginosa*. In contrast, the LD50 of the same product was determined on other earthworm species (Annelid) ;1262mg/kg on *Eisenia andrei* by Vermeulen et al (2001), 500 mg/kg on *Perionyx excavatus* by De Silva et al (2010).

By contrast, a 100% mortality rate could be observed after only 7 days exposure to dursban at a dose of 720 mg/kg or to the mixture of dursban ± mancozeb at 720/800 mg/kg. These results were similar to those obtained by Yesguer (2015) on Annelids

submitted to a 720 mg/kg of dursban alone, and Zaidi (2017) on *Octodrilus complanatus* (Annelid) exposed to the mixture dursban ± mancozeb at a concentration of 720/800 mg/kg. This high toxicity of Chlorpyrifos was also reported by Morgado et al.(2016) on the terrestrial isopod *Porcellionides pruinosus* (75% mortality) and Palma et al. (2008) on three aquatic species *Daphnia magna*, *Vibrio fischeri* and *Thamnocephalus platyurus*.

The mixture dursban ± mancozeb causes a lower (63 %) mortality than that observed for the dursban tested alone (69%). Similar observations were reported by Zaidi (2017) on earthworms (*Octodrilus complanatus*). On the other hand, some studies have shown that the effect of a mixture does not necessarily reflect the effects of individual substances (Padhi et al., 2008).

The possible interactions between the components of a mixture of pollutant are quite complex and make it very difficult to predict the overall effect (Lodivici et al., 1994). The toxic effects and significant mortality induced by dursban are explained by the high liposolubility of this insecticide; this property allows it to penetrate easily into the tissues of animals (Rice et al., 1997). After penetration inside the body, these products inhibit the functioning of the enzyme acetylcholine esterase (neurotransmitter of the nervous system) (Mileson et al.,1998; Venkateswara et al., 2004 ; Sanchez et al.,2014 ; Morgado et al.,2018) and thus prevent acetylcholine degradation (Regnault,

2005). This neurotransmitter accumulates in the synapses and causes hyperexcitation leading to death.

The acute toxicity test of dursban on *Armadillidium vulgare* revealed a lethal dose of 50% of the population after 14 days ($LD_{50} = 72.77$ mg/kg), this dose is close to that found by Hu et al. (2004) (83.626 mg/kg) and Zhou et al. (2007) (91.78 mg/kg) on earthworms (*Aporrectodea caliginosa*). Yesguer (2015) determined an LD_{50} of 60mg/kg with the same pesticide on *Aporrectodea caliginosa* (Annelid), whereas Lister et al. (2011) reported an LD_{50} between 26.6 and 38.9 mg/kg on *Lumbricus rebus*. Ma et al. (1993) and Chen et al. (2014) reported that dursban has a high toxicity on *Eisenia foetida* (Annelid) with respective values of 1077 mg/kg and 384.9 mg/kg. According to Kula et al. (1997), the variation in the LD_{50} value of the pesticide is due to the difference in substrate composition, environmental conditions and species tolerance. According to Belfroid et al. (1993); Kula et al. (1997); De silva et al. (2009) it varies with organic matter, texture, pH, temperature and exposure time. The difference observed between the LD_{50} of dursban in *Aporrectodea caliginosa* earthworms by Yesguer (2015) and that obtained on a terrestrial crustacean (*Armadillidium vulgare*), with the same environmental conditions and concentrations of this insecticide, is mainly due to the biological characteristics of the species studied. Like all crustaceans, *Armadillidium vulgare* is protected by a chitinous cuticle, while earthworms (Annelid) have a thin epidermis. Further -more, the diet of earthworms is composed of soil particles (Eggleton, 2006), making them highly vulnerable to contaminants compared to woodlice.

The modification of the structure of the cuticle observed in woodlice under the effect of pesticides made this cuticle fragile causing the destruction of the marsupium, which induces the release of mucus and the expulsion of eggs. The paralysis noticed in woodlice treated with the dursban is not a consequence of moulting since these individuals were not moulting during this period referring to Montesanto (2017). This paralysis has already been reported by Louat (2013) on the insect *Drosophila melanogaster* with the same insecticide (Chlorpyrifos). It is due to the inhibition of the enzyme Acetylcholinesterase which causes an accumulation of Acetylcholine in the cholinergic synapse leading to hyperexcitation of neurons. Paralysis is finally leading to death in insect (Louat, 2013).

CONCLUSION

The acute toxicity test carried out showed that dursban caused 100% mortality in the population at the dose used by farmers (720 mg/kg). The second pesticide mancozeb, caused less toxicity (19.5% mortality). When a mixture of the two pesticides was tested, the toxicity was lower (63% mortality) compared to the dursban alone. The contamination of woodlice by dursban also reveals an alteration in the marsupial cavity. The latter causes the eggs to be expelled outside the cavity hindering the embryonic development of

woodlice. Dursban, an insecticide widely used in agriculture, definitely represents a danger for all soil fauna. In the long term, the frequent use of dursban would be responsible for the decrease in the abundance of soil biodiversity reducing its biological activity. Consequently, this could lead to the loss of soil fertility, and agricultural yields.

ACKNOWLEDGEMENTS

We would like to thank the Editor-in-Chief and the anonymous reviewers who will review the first drafts of our manuscript for their valuable comments and suggestions. A big thank you to our proofreaders for the revision of English (Mrs Naziha Moumene, Djamilia Mezhoud). We would like to thank Mr. Ryad Moulai, director of our research laboratory (Laboratory of Applied Zoology and Animal Ecophysiology), for having provided us with all the necessary resources to carry out our research, as well as the engineer from the same laboratory (Habiba Baaziz) for his availability.

AUTHORS CONTRIBUTION

Moumene M.: Laboratory work; Sayah C. M.: Supervisor; Mouloud M. N., Kadji H. D.: Statistical study; Yesguer S.: Preparation of pesticide dilutions; Mahdeb M.: Soil analysis; Salhi A. M.: Field sampling; Charfi F.: Identification of species; Montesanto G., Habould C.: Language correction and final revision of the manuscript.

FUNDING

The authors declare that there were no significant financial support for this work that could have influenced its outcome.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

REFERENCES

- ASB, Agricultural Service of Bejaia, Algeria, 2017.
- ASISD, Agricultural Statistics and Information Systems Department, Ministry of Agriculture and Rural development and Fisheries, Algeria, 2017.
- Belfroid A, Seinen W, Van Gestel K, Hermens J, The acute toxicity of chlorobenzenes for earthworms (*Eisenia andrei*) in different exposure systems. *Chemosphere*. 26, 2265-2277, 1993.
- Bostanian N, Thistlewood H, Racette G, Effects of five fungicides used in Quebec apple orchards on *Amblyseius falia* Gies (Garman) (Phytoseiidae: Acari). *J Hort Sci Biotechnol*. 73(4) 527-530, 2015.
- Bureš S, Weidinger K, Sources and timing of calcium intake during reproduction in flycatchers. *Oecologia*. 137(4), 634- 641, 2003.
- Camila-Wood T, Zimmer M, Can terrestrial isopods (Isopoda :Oniscidea) make use of biodegradable plastics? *Appl. Soil Ecol*. 77, 72-79, 2014.
- Camila-Wood T, Kostanjč, Rok A, Paula B, Strus J, Morphology, microhabitat selection and life-

- history traits of two sympatric woodlice (Crustacea: Isopoda: Oniscidea): a comparative analysis. *Zool.Anz*,2017.
- Chen C, Yanhua W, Xueping Z, Qiang, W, Yong zhong Q, Comparative and combined acute toxicity of butachlor, imidacloprid and chlorpyrifos on earthworm, *Eisenia fetida*. *Chemosphere*.100 ,111-115,2014.
- Coline D, Effets des pesticides de la vigne sur le cycle biologique de l'escargot dans divers contextes d'exposition, Thèse de l'Université de Franche-Comté .Spécialité : Sciences de la Vie et de l'Environnement, École doctorale: Homme, Environnement, Santé.327,2011.
- Cortet J, Gomot-De Vaullery A., Poinsot Balagu er N,Gomot L, Texier C,The use of invertebrate soil fauna in monitoring pollutant effects. *Eur. J. Soil Biol.* 35 ,115-134,1999.
- Curry JA, New Program to Research Issues of Global Climate in the Arctic.*EOS.* 75 ,249-252,1994.
- De Silva Pathiratne A, Van G, Influence of temperature and soil type on the toxicity of three pesticides to *Eisenia andrei*. *Chemosphere.* 76,1410-1415,2009.
- De Silva Asoka P, Cornelis A, Toxicity of chlorpyrifos, carbofuran, mancozebe and their formulations to the tropical earthworm *Perionyx excavates*. *Appl Soil Ecol.*44, 56-60,2010.
- Eggleton P, The termite gut habitat: its evolution and co-evolution. In: intestinal microorganisms of soil invertebrates. König H &Varma A (eds), Springer-Verlag, Berlin. pp.373-403. 2006.
- Fabre R, Truhaut R, Toxicologie des Produits Phytopharmaceutiques. Société d'Édition d'Enseignement Supérieur - SEDES: Paris, pp.272,1954.
- Grosset-Souty C, BadenhauerI, Reynolds JD, Morel A, Investigations on the potential of woodlice as bioindicators of grassland habitat quality. *Eur. J. Soil Biol.*41,109-116,2005.
- Hassall M, Sutton, SL, The role of isopods as decomposers in a dune grassland ecosystem.*Scientific Proceedings of the Royal Dublin Society* A6. 1978, pp. 235-245, 1994.
- Hu Xiuqing Q, Cang T, Wu M, Study on toxicity and safety evaluation of chlorpyrifos and fenprothrin to earthworm (*Eisenia foelide*). *Pesticide Science and Administration.*25 , 10-11,2005.
- Ioriatti c, Pasqualini E, Toniolli A, Effects of the fungicides mancozebe and dithianon on mortality and reproduction of the predatory mite *Amblyseius Anderson*.*Exp Appl Acarol.*15 (2) , 109-116,2015.
- ISO, Qualité du sol, Effets des polluants vis-à-vis des vers de terre – Partie1 : Détermination de la toxicité aiguë vis à vis de *Eisenia fetida* / *Eisenia andrei*. Organisation internationale de normalisation, Norme internationale ISO/DIS 11268-1, 2011.
- Jänsch S, Garcia M, Römbke J, Acute and chronic isopod testing using tropical *Porcellionides pruinosus* and three model pesticides.*Eur. J. Soil Biol.*41,143-152,2018.
- Kammenga JE, Dallinger R, Donker MH, Köhler HR, Simonsen V, Triebkorn R, Weeks JM, Biomarkers in terrestrial invertebrates for ecotoxicological soil risk assessment.*Rev Environ Contam Toxicol.* 164 , 93-147, 2005.
- Koissi Savi M, Tchandao Mangamana E, Deguenon JM, Gbèmemali G, Hounmenou Glèlè Kakai R, Determination of Lethal Concentrations Using an R Software Function Integrating the Abbott Correction. *JASTA* 7, 25-30,2017.
- Kula H, Larink O, Development and standardization of test methods for the prediction of sublethal effects of chemicals on earthworms. *Soil Biol. Biochem.* 29, 635-639, 1997.
- Lister L, Svendsen C, Wright J, Hooper HL, Spurgeon DJ, Modelling the joint effects of a metal and a pesticide on reproduction and toxicokinetics in Lumbricid earthworms. *Environ Int.*37 ,663-670, 2011.
- Lodovici M, Aiolli S, Monserrat C, Dolara P, Medica A, Di Simplicio P, Effect of a mixture of 15 commonly used pesticides on DNA levels of 8-hydroxy-2-deoxyguanosine and xenobiotic metabolizing enzymes in rat liver. *J. Environ Pathol Toxicol Oncol.*13 (3) ,163-168, 1994.
- Louat F, Etude des effets liés à l'exposition aux insecticides chez un insecte modèle, *Drosophila melanogaster*.Thèse de Doctorat : Université d'Orléans.2013.
- Ma W, Bodt J, Differences in toxicity of the insecticide Chlorpyrifos to six species of earthworms (*Oligochaeta*, *Lumbricidae*) in standardized soil tests. *Bull. Environ.Contam.Toxicol.* 50 ,864-870, 1993.
- Mileson BE, Chambers JE, Chen WL, Dettbarn W, Ehrich M, Eldefrawi AT, aylor DW, Hamernik K, Hodgson E, Karczmar AG, Padilla S, Pope CN, Richardson RJ, Saunders DR, Sheets LP, Sultatos LG, WallaceKB, Common mechanism of toxicity: A case study of organophosphorus pesticides. *Toxicol Sci.* 41 ,8-20, 1998.
- Montesanto G, Cividini S, The moult cycle of the terrestrial isopod *Armadillo officinalis* Duméril, 1816 (Crustacea: Isopoda: Oniscidea).pp.1-11,2017.
- Morgado R, Pedro-Gomesa AD, Nuno Ferreira GC, Diogo-Caradosa N, Miguel Santosb JG, Mamadeu-Soaresa, M.V.,Loureiro, S. Toxicity interaction between chlorpyrifos, mancozeb and soil moisture to the terrestrial isopod *Porcellionides pruinosus*. *Chemosphere.* 144 , 1845-1853, 2008.
- Morgado R, Nuno Ferreira GC, Diogo-Cardoso N, Patrícia-Silva V, Amadeu MV, Soares M, Loureiro S, Joint effects of chlorpyrifos and mancozeb on the terrestrial isopod *Porcellionides pruinosus*: A multiple biomarker approach. *Environ Toxicol Chem.* 2018.
- OECD, Guideline for testing of chemicals."Earthworm, Acute Toxicity Tests. 1984.

- Padhi BK, Pelletier G, Williams A, Berndt-Weis L, Yauk C, Bowers WJ, Chu I, Gene expression profiling in rat cerebellum following in utero and lactational exposure to mixtures of methyl mercury, polychlorinated biphenyls and organochlorine pesticides. *Toxicological Letters*. 176 (2), 93-103, 2008.
- Palma P, Palma VL, Fernandes RM, Soares AMVM, Barbosa IR, Acute toxicity of atrazine, endosulfan sulphate and chlorpyrifos to *Vibrio fischeri*, *Thamnocephalus platyurus* and *Daphnia magna*, relative to their concentrations in surface waters from the Alentejo region of Portugal. *Bull Environ Contam Toxicol*. 81, 485-489, 2008.
- Pretty J. 2008. Agricultural sustainability: concepts, principles and evidence. *Phil. Trans. R. Soc. B363*, 447-465 10.1098/rstb.2007.2163.
- Regnault-Roger C, Enjeux phytosanitaires pour l'agriculture et l'environnement. *TEC & DOC*, 1013, 2005.
- Rice P, Drewes C, Klubertanz T, Bradbury S, Coats J, Acute toxicity and behavioral effects of Chlorpyrifos, permethrin, phenol, strychnine, and 2,4-dinitrophenol to 30-days-old Japanese Medaka (*Oryzias latipes*). *Setac Journals*. 16, 696-704, 1997.
- Saline G, Bio surveillance humaine, biomarqueurs et bio surveillance environnementale. *Annales Pharmaceutiques Française*. 70 : 199-203, 2012.
- Sanchez-Hernandez Juan C, Narvaez C, Sabat P, Martínez Mocillo S, Integrated biomarker analysis of chlorpyrifos metabolism and toxicity in the earthworm *Aporrectodea caliginosa*. *Sci Total Environ*. 490, 445-455, 2014.
- Szèpvölgyi j, Nagy k, SajgonèVukan k, Regoly-Mirei A, Soos K, Toth K, Pinter A, Antal M, Subacute toxicological examination of Dithane M-45. *Fd Chem. Toxic*. 27, 531-538, 1989.
- Van Gestel C A M, Loureiro S, idar P, Terrestrial isopods as model organisms in soil ecotoxicology: a review *Zookeys* (801): 127-162, 2018.
- Venkateswara Rao, J, Kavitha, P, Toxicity of azodrin on the morphology and acetylcholine esterase activity of the earthworm *Eisenia foetida*. *Environ. Res*. 96, 323- 327, 2004.
- Vermeulen LA, Reinecke AJ, Reinecke SA, Evaluation of the fungicide manganese zinc ethylene bis(dithiocarbamate) (mancozebe) for sublethal and acute toxicity of twenty-four insecticides to earthworm, *Eisenia foetida*, *Ecotox. Environ. Saf*. 48, 183-189, 2001.
- Viard- La Rocca B, Mise au point et validation, sur sites contaminés, (Etm-Hap) d'un test de bio surveillance en Microcosme : croissance et bioaccumulation par le gastéropode terrestre *Hélix Aspersa Aspersa*. These de doctorat Ecole Doctorale Sciences et Ingénierie des Ressources Procédés Produits Environnement Discipline : Sciences de la Vie Mention : Toxicologie de l'Environnement. 2004.
- Yesguer S, Evaluation de l'écotoxicité de certains pesticides sur les sols par l'utilisation d'un biotest : cas des lombricidés. Thèse de doctorat : Université de Bejaia. pp 31-40, 2015.
- Zaidi L, Safsaf A, Réponse éco physiologique des Lumbricidae à l'effet combiné de certains pesticides utilisés dans la Vallée de la Soummam, Master II en Bio ressources Animales et Biologie Intégrative. 2017.
- Zhou S, Duan C, Hui F, Chen Y, Wang X, Yu Z, Toxicity assessment for chlorpyrifos-contaminated soil with three different earthworm test methods. *J. Environ. Sci*. 19 (7), 854-858, 2007.