



## Abundance patterns of terrestrial isopods along an urbanization gradient

T. Magura<sup>1,4</sup>, E. Hornung<sup>2</sup> and B. Tóthmérész<sup>3</sup>

<sup>1</sup> Hortobágy National Park Directorate, H-4002 Debrecen, POB. 216, Hungary

<sup>2</sup> Department of Ecology, Szent István University, Faculty of Veterinary Science, Institute for Biology, H-1400 Budapest, POB 2, Hungary

<sup>3</sup> Department of Ecology, University of Debrecen, H-4010 Debrecen, POB. 71, Hungary

<sup>4</sup> Corresponding author. Hortobágy National Park Directorate, H-4002 Debrecen, POB. 216, Hungary.

Fax: +36 52 529940, e-mail: magura@hnp.hu

**Keywords:** GLOBENET, Habitat specialist hypothesis, Opportunistic species hypothesis, Urbanization.

**Abstract:** The abundance of terrestrial isopods (Isopoda: Oniscidea) was evaluated along an urban-suburban-rural gradient. We tested two hypotheses regarding the response of species: (i) habitat specialist hypothesis, according to which the abundance of the forest specialists would increase, while the abundance of the urban environment specialist isopods would decrease along the urban-rural gradient, and (ii) opportunistic species hypothesis (abundance of the generalist species would increase by increasing level of urbanization). The abundance of the forest specialist isopod *Trachelipus ratzeburgii* increased significantly along the studied gradient. An opposite tendency was observed for the abundance of the urban environment specialist isopod *Porcellio scaber*, as it was significantly higher in the urban area than in the suburban and rural sites. One generalist species (*Trachelipus rathkii*) gained dominance in the urban area, while other two generalists (*Armadillidium vulgare* and *Porcellium collicola*) showed no significant changes in abundance along the gradient.

**Nomenclature for isopods:** Schmalzfuss (2003).

### Introduction

Urbanization as a main form of anthropogenic activities is an increasingly important force shaping the landscape by creating patchworks of modified land types. Urbanization associated with changes in the temperature (Hawkins et al. 2004), pollution (Conti et al. 2004) and the area and the quality of natural habitat (Bolger et al. 2000, McIntyre et al. 2001, Gibb and Hochuli 2002) results in a densely populated, built-up, developed and often highly disturbed urban area (city centre) which is surrounded by areas of decreasing development and habitation with moderate (suburban area) or light disturbance (rural area) level (Dickinson 1996). Such urban-rural gradients representing decreasing intensity of human influence exhibit similar patterns throughout the world. Despite the prevalence and acceleration of urbanization and that urbanization is considered as one of the primary causes for biodiversity loss, little is known on whether or not changes caused by urbanization affect biodiversity in similar way across the globe (Niemelä et al. 2000).

In 1998, an international research project, the "Globenet" project (Global Network for Monitoring Landscape Change), to search for generalizations in urbanization impacts on biodiversity was initiated (Niemelä and Kotze 2000, Niemelä et al. 2000). As the effects of urbanization on biodiversity can be explored most effectively through investigations along urban-to-rural gradients (McDonnell et al. 1997), the Globenet

project also employs this gradient approach. During field survey, the Globenet project uses a common, standardized methodology (pitfall trapping) for ground dwelling arthropods (like carabid beetles, spiders and terrestrial isopods). In the frame of the Globenet project, several papers investigating carabid assemblages were published (Alarukka et al. 2002, Niemelä et al. 2002, Ishitani et al. 2003, Venn et al. 2003, Magura et al. 2004, 2005, 2006, 2008, Gaublumme et al. 2005, Tóthmérész and Magura 2005, Sadler et al. 2006, Elek and Lövei 2007). Studies analyzing other target arthropod assemblages along the urbanization gradient are very limited (for spiders: Alarukka et al. 2002). Terrestrial isopods (Isopoda: Oniscidea) are considered reliable biological indicator organisms of environmental stress both at the assemblage and species level (Dallinger et al. 1992, Jones and Hopkin 1998, Paoletti and Hassall 1999). Therefore, terrestrial isopods are potential targets for studying the effect of urbanization.

Traditionally, urbanization is considered as a form of disturbance (Rebele 1994). Several hypotheses were put forward to explain the effects of disturbance on biotic communities. Species with different ecological characteristics may respond differently to urbanization (Magura et al. 2004): forest specialist species may suffer, while species associated with the altered urban habitat may benefit from the disturbance and habitat alteration caused by urbanization. Therefore, analyzing the overall species abundance along the gra-

dient may be misleading and may disguise basic ecological rules (Magura et al. 2004). In an earlier paper (Hornung et al. 2007), we tested the Intermediate Disturbance Hypothesis and the Increasing Disturbance Hypothesis on isopods; trends in the total number of individuals, number of species, and three diversity indices (Shannon, Simpson, and Berger-Parker) were evaluated. Hornung et al. (2007) did not find significant differences in these parameters between the studied sites (urban, suburban, and rural). These findings gave rise to the idea that groups of species with different ecological characteristics (forest specialists, urban environment specialists, generalists) should be tested as their reactions to urbanization could be different and could cancel each other out when evaluated together. Since increasing disturbance by urbanization affects primarily the habitat specialist species, we hypothesized that abundance of the forest specialist isopods should increase, while abundance of the urban environment specialist isopods (or synanthropic species) should decrease from the urban area to the rural one (habitat specialist hypothesis, Magura et al. 2004). Moreover, we hypothesized that the abundance of the generalist species should be higher in the urban area, as predicted by the opportunistic species hypothesis (Gray 1989).

## Material and methods

### Study area and sampling design

Following the Globenet sampling protocol, terrestrial isopods were studied along an urban-suburban-rural gradient in Debrecen (Eastern-Hungary), the second largest city of the country (Magura et al. 2004). The urban, suburban and rural sampling areas were situated in once continuous patches of old forest (Convallario-Quercetum forest association; 100 yrs) dominated by English oak (*Quercus robur*). These forest patches covered an area of at least 6 ha. The distinction of sampling areas (urban, suburban and rural) was based on the ratio of built-up area to natural habitats measured by the Arc-View GIS program using an aerial photograph in a square of 1 km<sup>2</sup> size around the sampling area. Buildings, roads and asphalt covered paths were regarded as built-up area. In the urban area, the built-up part exceeded 60%, in the suburban area it was approximately 30%, while in the rural one there was no built-up area. The forest patches in the urban area belong to an urban park with several asphalt covered paths and strongly thinned shrub layer. In the suburban area fallen trees were removed, while in the rural area forest management was only occasional at a low-intensity level. Distances between the sampling areas (urban, suburban, rural) were at least 1-1 km, as prescribed by the general methodology of the Globenet project (Niemelä et al. 2000). Four sites, at least 50 m from each other (in order to achieve independency, see Digweed et al. 1995), were selected within each sampling area. Terrestrial isopods were collected at each of the 4 sites of the 3 sampling areas using unbaited pitfall traps, consisting of plastic cups (diameter 65 mm, volume 250 ml) containing 75% ethylene glycol as a killing-preserving solution from the end of March to the end of November, 2001. Traps

were emptied fortnightly. Ten traps were placed randomly at least 10 m apart at each site. This resulted in a total of 120 traps scattered along the urban-rural gradient (3 area × 4 sites × 10 traps). Each pitfall trap was at least 50 m from the nearest forest edge, in order to avoid edge effects (Molnár et al. 2001). The traps were covered with bark pieces to protect them from litter and rain (Spence and Niemelä 1994). For analysis, catches from each trap were pooled for the whole year. Of course, other sampling methods (e.g. hand sorting, Tullgren funnel extraction, litter sieving) could enhance the possibility to catch rare and/or small sized, soil dwelling species.

### Data analyses

To test differences in the overall isopod abundance, and in the abundance of the trapped isopod species among the three sampling areas (urban, suburban and rural) and among the 12 sites, nested analyses of variance (ANOVA) were performed. Data of individual traps were used (sites nested within the sampling areas). Normal distribution of the data was achieved by log(x+1) transformation (Sokal and Rohlf 1995). Ecological characteristics (forest specialists, urban environment specialists, generalists) of the terrestrial isopod species caught were based on the literature (Schmalfluss 2003). The urban environment specialist species are frequently non-native, but mainly invasive and/or established introduced ones (Vilisics et al. 2007b). When ANOVA revealed a significant difference between the means, the Tukey test was performed for multiple comparisons among means.

## Results


The total isopod catch consisted of 9115 individuals representing 6 species. 3548 individuals belonging to 6 species were captured in the urban, 5 species and 2720 individuals in the suburban, and 4 species and 2847 individuals in the rural area. The most abundant species was *Armadillidium vulgare*, which made up 72% of the total catch. Out of the trapped 6 species, there were three generalist species (*Armadillidium vulgare*, *Porcellium collicola* and *Trachelipus rathkii*), one forest specialist (*Trachelipus ratzeburgii*) and two urban environment specialists, as established introduced species (*Cylisticus convexus* and *Porcellio scaber*) (Table 1).

Analyzing the trap-level data by nested ANOVA we found that there were no statistically significant differences in the overall abundance of isopods across the urban-rural gradient ( $F=0.4859$ ; d.f.=2, 9;  $p>0.05$ ). The abundance of the forest specialist isopod *Trachelipus ratzeburgii* increased significantly from the urban area toward the rural one ( $F=14.3469$ ; d.f.=2, 9;  $p<0.01$ ; Fig. 1A).

An opposite tendency was observed regarding the abundance of *Porcellio scaber* (urban environment specialist isopod), which was significantly higher in the urban area than in the suburban and rural ones ( $F=6.1014$ ; d.f.=2, 9;  $p<0.05$ ; Fig. 1B). *Cylisticus convexus*, the other urban envi-

**Table 1.** The catches of the terrestrial isopod species and their ecological category along an urban-rural gradient.

| Species  | Ecological category          | Urban | Suburban | Rural | Total |
|--|------------------------------|-------|----------|-------|-------|
| <i>Armadillidium vulgare</i> (Latreille, 1804) | generalist                   | 2088  | 2280     | 2218  | 6586  |
| <i>Cylisticus convexus</i> (De Geer, 1778)     | urban environment specialist | 1     | 2        | 0     | 3     |
| <i>Porcellium collicola</i> (Verhoeff, 1907)   | generalist                   | 272   | 245      | 226   | 743   |
| <i>Porcellio scaber</i> Latreille, 1804        | urban environment specialist | 28    | 0        | 0     | 28    |
| <i>Trachelipus rathkii</i> (Brandt, 1833)      | generalist                   | 1143  | 10       | 64    | 1217  |
| <i>Trachelipus ratzeburgii</i> (Brandt, 1833)  | forest specialist            | 16    | 183      | 339   | 538   |



**Figure 1.** Average abundance values ( $\pm$ SE) of *Trachelipus ratzeburgii*, a forest specialist species (A), *Porcellio scaber*, an urban environment specialist species (B) and *Trachelipus rathkii*, a generalist species (C) calculated for the pitfall traps along an urban-suburban-rural gradient. Letters a, b and c indicate significant ( $p < 0.05$ ) differences based on the Tukey multiple comparison test.

ronment specialist species was excluded from the analysis because of its low catches (only 3 individuals were caught).

The abundance of *Trachelipus rathkii*, a generalist species, was significantly higher in the urban area compared to the suburban and rural ones ( $F=9.4200$ ; d.f.=2, 9;  $p<0.01$ ; Fig. 1C). The other two generalist species (*Armadillidium vulgare* and *Porcellium collicola*) showed no significant changes in abundance along the gradient ( $F=0.3632$ ; d.f.=2, 9;  $p<0.05$ ; and  $F=0.5999$ ; d.f.=2, 9;  $p>0.05$ ; respectively).

**Discussion**

*Overall isopod abundance*

Several previous papers have emphasized that terrestrial isopods respond predictably to disturbance and that disturbance causing habitat alteration impacted assemblage composition, leading to significant changes in the abundance and species richness (Kalisz and Powell 2004, Pitzalis et al. 2005, Tsukamoto and Sabang 2005). In contrast, we did not find

any significant difference in the abundance of isopods across the city wide gradient.

The lack of significant difference in the overall abundance may be caused by the dissimilar response of isopods with different habitat affinity to disturbance. Forest specialists may suffer, while generalists and urban environment specialist may benefit from the disturbance and habitat alteration caused by urbanization (Magura et al. 2004).

#### *Abundance of forest specialist, urban environment specialist and generalist isopods*

Contrary to the overall terrestrial isopod assemblages, the forest specialist and the urban environment specialist species responded significantly to the changes in habitat conditions caused by the urbanization. Our findings illustrated that forest specialist species suffer, while urban environment specialist species benefit from the disturbance and habitat alteration. Our results indicate that urbanization has a very strong effect on the forest specialist isopod species *Trachelipus ratzeburgii*. This forest species is more stenotopic, demanding microsites with favorable microclimate, the presence of dead and decaying trees, and significant cover of coarse woody debris, leaf litter, shrubs and herbs, together forming an undisturbed habitat (Korsós et al. 2002). Habitat alteration caused by urbanization appears to eliminate favorable microsites for forest species and therefore contributes to their decline of abundance at the disturbed areas. Along the studied urbanization gradient, disturbance was the highest in the urban area (paved paths, thinned shrub layer), it was moderate in the suburban area (fallen trees removed), and was lowest in the rural one. This decreasing disturbance was also expressed by the increasing abundance of the forest specialist isopod species. Judas and Hauser (1998) studying the distribution patterns of isopods in beech forests emphasized that *Trachelipus ratzeburgii* was the most abundant in the habitat patches with dead wood material. Jabin et al. (2004) also showed that coarse woody debris was a significant positive predictor for the abundance of the terrestrial isopod species in a forested habitat. In the same forest patches sampled in this study, Hornung et al. (2007) evaluated the relationships between the abundance of isopod species and environmental factors. Their results proved that the abundance of *Trachelipus ratzeburgii* increased with the amount of the decaying wood material. Moreover, there was negative correlation between the abundance of *Trachelipus ratzeburgii* and the ground and air temperature indicating that this species prefers closely natural habitats, which are usually characterized by lower ground and air temperature (Hornung et al. 2007).

Our findings show that the urban park is the most favorable habitat for the urban environment specialist *Porcellio scaber*. Its abundance was significantly higher there compared to the suburban and rural area. The exclusive appearance of *Porcellio scaber* in the urban park is in accordance with its habitat preference. Contrary to some other European countries with Atlantic climate influence, in Hungary this species can be found only in human settlements, in and

around houses because of the heat island effect and moist shelters (Schmalfuss 2003). The preference of *Porcellio scaber* for higher temperatures was shown by Hornung et al. (2007), as its abundance increased as the ground and the air temperature on the surface increased. Moreover, this urban environment specialist species may tolerate heavy metals originated from air pollution by accumulating them in vesicles in the hepatopancreas (Paoletti and Hassall 1999). Thus, they can also survive in the polluted habitats, although their body size can decline significantly (Jones and Hopkin 1998).

Our result partly supported the prediction of the opportunistic species hypothesis, as one of the generalist species, *Trachelipus rathkii* gained dominance in the disturbed urban area. Hornung et al. (2007) showed that the generalist *Trachelipus rathkii* preferred the urban site, which usually can be characterized by higher average temperature (heat islands effect). The other two generalist species (*Armadillidium vulgare* and *Porcellium collicola*), however, showed no significant changes in abundance along the gradient. These facts indicate that there is no clear, unique pattern along the urban-rural gradient for generalist species, because their abundance pattern is controlled by their autecological characteristics in complex interaction with the environmental variables, and urbanization/disturbance level. Nevertheless, previous studies on ground beetles (Niemelä et al. 2002 for the Canadian and Finnish Globenet sites, Ishitani et al. 2003, Magura et al. 2004, Sadler et al. 2006, Elek and Lövei 2007) and on ants (Vepsäläinen et al. 2008) confirmed the opportunistic species hypothesis.

#### *Urban forests and biodiversity*

We trapped six terrestrial isopod species in the studied urban park. This species richness is in accordance with the overall isopod diversity of natural forests in Hungary where the average isopod species richness in natural or semi-natural deciduous forests is around 5-6 (Loksa 1966). However, the present study also stressed that habitat modification caused by urbanization altered remarkably the terrestrial isopod assemblages, as the abundance of the forest specialist isopods decreased, while that of the urban environment specialist isopods and partly generalist isopods increased with the increasing disturbance level. Thus, urbanization could be one of the leading reasons of alteration in indigenous arthropod assemblages as well (Davis 1978). On the other hand, urban parks and other urban green areas created by the rising urbanization have a vital recreational importance and increase the quality of urban life. Therefore, there is a growing need for appropriate management strategies which consider simultaneously recreational, economic and conservation criteria (Gilbert 1989). We propose that extensive modification of habitats should be avoided, as these alterations are accompanied by unfavorable changes in the microclimatic, abiotic and biotic conditions of the area. Positive effects of soft management (the cut grass, plant material, trimmed branches and litter were returned to the urban forest patches) on isopod di-

versity and abundance was proved in an urban park of Sorø, Denmark (Vilisics et al. 2007a).

**Acknowledgments.** Research was supported by the grants of the Hungarian Scientific Research Fund (OTKA research grant nos. F61651 (to TM) and T043508 (to EH)). TM is grateful for the support of Bolyai Fellowship. We wish to express our thanks to T. Molnár, F. Vilisics and Z. Elek for their help in field and laboratory works.

## References

- Alaruikka, D.M., D.J. Kotze, K. Matveinen and J. Niemelä. 2002. Carabid and spider assemblages along an urban to rural gradient in Southern Finland. *J. Insect Conserv.* 6: 195-206.
- Bolger, D.T., A.V. Suarez, K.R. Crooks, S.A. Morrison and T.J. Case. 2000. Arthropods in urban habitat fragments in southern California: Area, age, and edge effects. *Ecol. Appl.* 10: 1230-1248.
- Conti, M.E., M. Tudino, J. Stripeikis and G. Cecchetti. 2004. Heavy metal accumulation in the lichen *Evernia prunastri* transplanted at urban, rural and industrial sites in central Italy. *J. Atmospheric Chem.* 49: 83-94.
- Dallinger, R., B. Berger and S. Birkel. 1992. Terrestrial Isopods - useful biological indicators of urban metal pollution. *Oecologia* 89: 32-41.
- Davis, B.N.K. 1978. Urbanisation and the diversity of insects. In: L.A.Mound and N. Waloff (eds), *Diversity of Insect Faunas*. Blackwell, London. pp. 126-138.
- Dickinson, R.E. 1996. The process of urbanization. In: F.F. Darling and J.P. Milton (eds), *Future Environments of North America*. Natural History Press Garden City. pp. 463-478.
- Digweed, S.C., C.R. Currie, H.A. Cárcamo, and J.R. Spence. 1995. Digging out the digging-in effect of pitfall traps: influences of depletion and disturbance on catches of ground beetles (Coleoptera: Carabidae). *Pedobiol.* 39: 561-576.
- Elek, Z. and G.L. Lövei. 2007. Patterns in ground beetle (Coleoptera: Carabidae) assemblages along an urbanisation gradient in Denmark. *Acta Oecologica* 32: 104-111.
- Gaublomme, E., H. Dhuyvetter, P. Verdyck and K. Desender. 2005. Effects of urbanisation on carabid beetles in old beech forests. *DIAS Report* 114: 111-123.
- Gibb, H. and D.F. Hochuli. 2002. Habitat fragmentation in an urban environment: large and small fragments support different arthropod assemblages. *Biol. Conserv.* 106: 91-100.
- Gilbert, O.L. 1989. *The Ecology of Urban Habitats*. Chapman and Hall, London.
- Gray, J.S. 1989. Effects of environmental stress on species rich assemblages. *Biol. J. Linnean Soc.* 37: 19-32.
- Hawkins, T.W., A.J. Brazel, W.L. Stefanov, W. Bigler and E.M. Safell. 2004. The role of rural variability in urban heat island determination for Phoenix, Arizona. *J. Appl. Meteorol.* 43: 476-486.
- Hornung E., B. Tóthmérész, T. Magura and F. Vilisics. 2007. Changes of isopod assemblages along an urban-suburban-rural gradient in Hungary. *Eur. J. Soil Biol.* 43: 158-165.
- Ishitani M., D.J. Kotze and J. Niemelä. 2003. Changes in carabid beetle assemblages across an urban-rural gradient in Japan. *Ecography* 26: 481-489.
- Jabin M., D. Mohr, H. Kappes and W. Topp 2004. Influence of dead-wood on density of soil macro-arthropods in a managed oak-beech forest. *Forest Ecol Manage.* 194: 61-69.
- Jones, D.T. and S.P. Hopkin. 1998. Reduced survival and body size in the terrestrial isopod *Porcellio scaber* from a metal-polluted environment. *Environmental Pollution* 99: 215-223.
- Judas M. and H. Hauser. 1998. Patterns of isopod distribution: From small to large scale. *Israel J. Zool.* 44: 333-343.
- Kalisz, P.J. and J.E. Powel. 2004. Exotic isopods (Crustacea: Isopoda) in disturbed and undisturbed forest soils on the Cumberland Plateau of Kentucky, USA. *Natural Areas Journal* 24: 54-56.
- Korsós, Z., E. Hornung, K. Szlávecz and J. Kontschán. 2002. Isopoda and Diplopoda of urban habitats: New data to the fauna of Budapest. *Annales Historico-Naturales Musei Nationalis Hungarici* 94: 45-51.
- Loksa, I. 1966. *Die Bodenzooökologischen Verhältnisse der Flaumeichen-Buschwälder Südostmitteleuropas*. Akadémiai Kiadó, Budapest.
- Magura, T., B. Tóthmérész and G.L. Lövei. 2006. Body size inequality of carabids along an urbanisation gradient. *Basic and Appl. Ecol.* 7: 472-482.
- Magura, T., B. Tóthmérész and T. Molnár. 2004. Changes in carabid beetle assemblages along an urbanisation gradient in the city of Debrecen, Hungary. *Landsc. Ecol.* 19: 747-759.
- Magura, T., B. Tóthmérész and T. Molnár. 2005. Species richness of carabids along a forested urban-rural gradient in eastern Hungary. *DIAS Report* 114: 209-217.
- Magura, T., B. Tóthmérész and T. Molnár. 2008. A species-level comparison of occurrence patterns in carabids along an urbanisation gradient. *Landsc. Urban Planning* 86: 134-140.
- McDonnell, M.J., S.T.A. Pickett, P. Groffman, P. Bohlen, R.V. Pouyat, W.C. Zipperer, R.W. Parmelee, M.M. Carreiro and K. Medley. 1997. Ecosystem processes along an urban-to-rural gradient. *Urban Ecosystems* 1: 21-36.
- McIntyre, N.E., J. Rango, W.F. Fagan and S.H. Faeth. 2001. Ground arthropod community structure in a heterogeneous urban environment. *Landsc. Urban Planning* 52: 257-274.
- Molnár, T., T. Magura, B. Tóthmérész and Z. Elek. 2001. Ground beetles (Carabidae) and edge effect in oak-hornbeam forest and grassland transects. *Eur. J. Soil Biol.* 37: 297-300.
- Niemelä, J. and D.J. Kotze. 2000. GLOBENET: the search for common anthropogenic impacts on biodiversity using carabids. In: P. Brandmayr, G.L. Lövei, T.Z. Brandmayr, A. Casale and A.V. Taglianti (eds), *Natural History and Applied Ecology of Carabid Beetles*. Pensoft Publisher, Sofia, Bulgaria. pp. 241-246.
- Niemelä, J., J. Kotze, A. Ashworth, P. Brandmayr, K. Desender, T. New, L. Penev, M. Samways and J. Spence. 2000. The search for common anthropogenic impacts on biodiversity: a global network. *J. Insect Conserv.* 4: 3-9.
- Niemelä, J., J. D. Kotze, S. Venn, L. Penev, I. Stoyanov, J. Spence, D. Hartley and E. Montes de Oca. 2002. Carabid beetle assemblages (Coleoptera, Carabidae) across urban-rural gradients: an international comparison. *Landsc. Ecol.* 17: 387-401.
- Paoletti, M.G. and M. Hassall. 1999. Woodlice (Isopoda: Oniscidea): their potential for assessing sustainability and use as bioindicators. *Agriculture Ecosystems and Environment* 74: 157-165.
- Pitzalis, M., S. Fattorini, E. Trucchi and M.A. Bologna. 2005. Comparative analysis of species diversity of Isopoda Oniscidea and

- Collembola communities in burnt and unburnt habitats in central Italy. *Ital. J. Zool.* 72: 127-140.
- Rebele, F. 1994. Urban ecology and special features of urban ecosystems. *Global Ecol. Biogeogr.* 4: 173-187.
- Sadler, J.P., E.C. Small, H. Fiszpan, M.G. Telfer and J. Niemelä. 2006. Investigating environmental variation and landscape characteristics of an urban-rural gradient using woodland carabid assemblages. *J. Biogeogr.* 33: 1126-1138.
- Schmalfuss, H. 2003. World catalog of terrestrial isopods (Isopoda: Oniscidea). *Stuttgarter Beiträge zur Naturkunde Serie A* 654: 1-341.
- Sokal, R.R. and F.J. Rohlf. 1995. *Biometry*. Freeman, New York.
- Spence, J.R. and J. Niemelä. 1994. Sampling carabid assemblages with pitfall traps: the madness and the method. *The Canadian Entomol.* 126: 881-894.
- Tsukamoto, J. and J. Sabang. 2005. Soil macro-fauna in an *Acacia mangium* plantation in comparison to that in a primary mixed dipterocarp forest in the lowlands of Sarawak, Malaysia. *Pedobiol.* 49: 69-80.
- Tóthmérész, B. and T. Magura. 2005. Affinity indices for environmental assessment using carabids. *DIAS Report* 114: 345-352.
- Venn, S.J., D.J. Kotze and J. Niemelä. 2003. Urbanization effects on carabid diversity in boreal forests. *Eur. J. Entomol.* 100: 73-80.
- Vepsäläinen, K., H. Ikonen and M.J. Koivula. 2008. The structure of ant assemblages in an urban area of Helsinki, southern Finland. *Ann. Zool. Fenn.* 45: 109-127.
- Vilisics, F., Z. Elek, G.L. Lövei and E. Hornung. 2007a. Composition of terrestrial isopod assemblages along an urbanisation gradient in Denmark. *Pedobiol.* 51: 45-53.
- Vilisics, F., P. Sóllymos and E. Hornung. 2007b. A preliminary study on habitat features and associated terrestrial isopod species. In: K. Tajovsky, J. Schlaghamersky and V. Pizl (eds), *Contributions to Soil Zoology in Central Europe II*. Institute of Soil Biology, Biology Centre, Academy of Sci. of the Czech Republic, České Budějovice. pp. 195-199.

Received May 13, 2008  
Revised June 20, 2008  
Accepted June 25, 2008