

Bug cemetery: a case study of terrestrial isopod distribution on a brick wall in the Czech Republic

Ivan Hadrián Tuf¹, Nelly Weissová¹

¹ Department of Ecology and Environmental Sciences, Palacký University, Olomouc, Czech Republic

Corresponding author: Ivan Hadrián Tuf (ivan.tuf@upol.cz)

Academic editor: Pallieter De Smedt | Received 4 October 2021 | Accepted 14 January 2022 | Published 18 May 2022

<http://zoobank.org/1AE60064-B993-4247-BD98-A52DD768EDE9>

Citation: Tuf IH, Weissová N (2022) Bug cemetery: a case study of terrestrial isopod distribution on a brick wall in the Czech Republic. In: De Smedt P, Taiti S, Sfenthourakis S, Campos-Filho IS (Eds) Facets of terrestrial isopod biology. ZooKeys 1101: 191–201. <https://doi.org/10.3897/zookeys.1101.76132>

Abstract

Although terrestrial isopods (Oniscidea) are primarily soil- and surface-dwelling invertebrates, they can also be found on tree trunks and walls. This study evaluated distribution patterns of terrestrial isopods on a brick wall during the first hours of night in autumn. Four species of terrestrial isopods were recorded with *Armadillidium versicolor* being the dominant one. Terrestrial isopods were distributed from ground level up to a height of 2 m, but preferred a 70–80 cm height band. The highest number of active individuals was observed 3 h after astronomical dusk. Potential predators of terrestrial isopods were abundant during the same time and at the same height.

Keywords

Isopoda, Oniscidea, synanthropic habitat, vertical distribution, woodlice

Introduction

Terrestrial isopods (Crustacea, Isopoda, Oniscidea) inhabit soil generally and feed on dead and rotting organic matter. They can be found in upper soil layers including the litter layer (Cole 1946). They shelter under bark, logs and stones or they are true troglobionts. There are several species climbing on vegetation as reported from tropical areas, such as Central America (e.g., Van Name 1936), Central (Schmidt 1999) and South Africa (Glazier and Kleynhans 2015) or St. Helena Island (Dutton and Pryce

2018). In Europe, there are only a few anecdotal reports about the presence of terrestrial isopods on trees, walls, and inside tree hollows for species of the genus *Armadillidium* (e.g., *A. pictum* Brandt, 1833, *A. depressum* Brandt, 1833, *A. pulchellum* (Zenker, 1798), *A. vulgare* (Latreille, 1804)), the genus *Porcellio* (*P. scaber* Latreille, 1804, *P. spinicornis* Say, 1818), *Oniscus asellus* Linnaeus, 1758, and occasionally other species (Růžicka et al. 1991; Alexander 2008, 2011; Gregory 2009; Božanić et al. 2013; Tracz 2013; Boeraeve et al. 2021). This above-ground activity is sometimes related to heavy rains (Abbott 1918; Standen 1921; Cole 1946) or spring floods (Fig. 1). However, a recent systematic inventory of arthropods in oak canopies in Norway revealed five species of terrestrial isopods with high prevalence of *A. pictum* (Thunes et al. 2021). Sutton (1972) noted the presence of *P. scaber* on tree trunks and brick walls feeding on green algae.

Although the nocturnal presence of *P. scaber* on trees is commonly known (Sutton 1972; Cloudsley-Thompson 1977; Warburg 1993), there are only a few systematic studies. The first one was by Brereton (1957), who systematically searched for terrestrial isopods on tree trunks near Oxford for two years; he reported four species found on tree trunk bases but only two of them (*P. scaber* and *O. asellus*) also at eye-level height (~ 1.8 m). During the year, isopods were the most numerous on trees in spring and in September/October. Brereton (1957) confirmed that *P. scaber* overwinters in moss pillows at the base of the trunk and the species climbs on branches during late summer: he found them in high numbers at 5.5 m height in early September. High abundance of terrestrial isopods in moss on the base of tree trunks were caused by downward migration of isopods of the canopy and the trunk, as confirmed with the use of trunk traps (Brereton 1957). The surprising opposite pattern of seasonal migration on trees for overwintering of *P. scaber* was suggested by Fritsche (1934; Vandel 1962). Prevailing descent migration from trees during a year was revealed for *Spherillo* spp. and *Trichoniscus* spp. in New Zealand (Moeed and Meads 1983), indicating breeding in canopies.

Another exhaustive study of *P. scaber* living on trees near Den Haag, the Netherlands, was published by Den Boer (1961), who studied its activity in the same years as Brereton (1957), 1953–1954 and 1953–1955, respectively. He described parameters of daytime shelters on the bark for *P. scaber*. He found that very few isopods, active during the night on tree trunks with such shelters, travel down to hide in the litter layer during daytime. Another interesting finding (recorded using the capture-mark-recapture method) is that although isopods are present on the bark during the whole night, activity of each individual isopod spans only approximately one hour (Den Boer 1961).

Sixty years later, aspects of distribution of terrestrial isopods on tree trunks were presented at the 11th International Symposium on Terrestrial Isopod Biology by Mechthold et al. (2021). They highlighted phenological differences between tree trunk active populations of *P. scaber* and *O. asellus* during the year.

Beside these studies on the activity of terrestrial isopods on tree trunks, there is only one paper studying distribution of isopods on walls. Cloudsley-Thompson (1958) counted terrestrial isopods on a stone wall at night and found that their numbers decrease with increasing wind speed. His explanation was that wind inhibits their activity due to a reduction of air humidity.



Figure 1. Springtails and terrestrial isopods climbing on tree trunks to avoid drowning during a spring flood. *Trachelipus rathkii* (Brandt, 1833), indicated by pink arrow, *Protracheoniscus politus* (C. L. Koch, 1841), indicated by orange arrow, and *Porcellium conspersum* (C. L. Koch, 1841), indicated by red arrow, are visible. Litovelské Pomoraví PLA (Czech Republic), 30 March 2006 (photographs IHT).

We also received anecdotal observations of terrestrial isopods climbing on walls of buildings during the night. In this study we present data on the distribution of terrestrial isopods on a brick wall in the Czech Republic to study (1) whether there is a temporal pattern in the distribution of particular species of terrestrial isopods, (2) whether its distribution depends on the air temperature or humidity, and (3) whether the distribution of predators corresponds with the distribution of terrestrial isopods.

Materials and methods

After a short pilot survey, we selected a study wall on which we found a high number of active terrestrial isopods during the night. This particular brick wall was found on the outskirts of the town of Kostelec na Hané (Czech Republic) at the local cemetery (49°31'06.0"N, 17°03'44.6"E). The length of the wall is 190 m and its height is ~ 2.5 m. The first 4 m of the wall is plastered, the rest are bare bricks standing on a 50 cm high stone foundation (Fig. 2A). Terrestrial isopods and their potential predators were studied along a 30 m long transect of the non-plastered part at the outward side of the wall (exposed to the east, i.e., influenced by the prevailing south-eastern wind) from the base to a height of 2 m. The wall was bordered by a freshly ploughed agricultural field and generally surrounded by an urban and agricultural landscape. The nearest forest is located at ~ 2 km. The nearest street lamp was located ~ 20 m away, but the studied side of the wall was not illuminated by artificial light.



Figure 2. **A** the studied brick cemetery wall **B** *Armadillidium versicolor* climbing on the wall (photographs NW).

Following Brereton (1957), the month of October (2019) was chosen as the best time to study terrestrial isopod activity. Previous observations indicated the highest abundance of terrestrial isopods between 21:00–23:00 h (i.e., 1.5–3.5 h after astronomical dusk which was at 19:38 on the first observation day; Fig. 2B). Therefore, terrestrial isopods and other invertebrates were observed daily from 20 to 26 October. Each daily observation started at $\sim 21:15$ (± 15 min) and lasted 90 min. At the start of each observation, we measured air humidity, air temperature, and light intensity (lux). For each observed individual, we noted species identity, time of observation, and height above the ground with a Dictaphone. Numbers of predatory invertebrates, i.e., centipedes, spiders, and harvestmen, were also noted to evaluate the pattern of distribution of potential predators. Observations were made using a hand torch with white light, since we were not interested in the behaviour of the animals, but only their position. No fleeing reaction was noted during the research. No individuals were collected during the observations. A few individuals for identification were taken a few days before the start of the first observation.

All data was entered into MS Excel, and we used CANOCO 5 (Šmilauer and Lepš 2014) for statistical analysis. Numbers of individuals of the different terrestrial isopod species were used as species variables. As environmental variables we used height (cm), air humidity (%), air temperature ($^{\circ}\text{C}$), time after astronomical dusk (minutes), and numbers of individuals of predators (summed numbers of centipedes (Chilopoda), spiders (Araneae), and harvestmen (Opiliones)). We used a redundancy analysis (RDA) for visualizing the relationship between number of terrestrial isopods




Figure 3. Number of invertebrates recorded per day on the wall during a 90-min observation during October 2019.

per species and environmental variables. We did not incorporate the effect of light intensity in the analysis, because measured values were very low and varied between 0 and 2 lx. Environmental variables that significantly explained variation in the terrestrial isopod distribution were used to calculate predictive GAM models (normal distribution of data).

Results

In total, 1221 terrestrial isopods belonging to four species were observed. By far the most numerous was *Armadillidium versicolor* Stein, 1859 (1020 individuals), followed by *Porcellio spinicornis* (112 ind.), *Armadillidium vulgare* (85 ind.), and *Porcellionides pruinosus* (Brandt, 1833) (4 ind.). Altogether, 266 spiders and only two centipedes and nine harvestmen were observed. The number of observed terrestrial isopods and predators decreased during the sampling period (Fig. 3), which could be due to the decreasing recorded temperature. However, this correlation was not significant (Pearson's $r = 0.37$ and $p = 0.414$ for temperature and number of observed terrestrial isopods, Pearson's $r = 0.48$ and $p = 0.276$ for temperature and number of observed predators). The highest mean numbers of individuals for (almost) all species were recorded at a height of 70–80 cm (Fig. 4).

The RDA for predicting the distribution of terrestrial isopods using environmental variables (Fig. 5) was statistically significant (pseudo-F = 28.3, $p = 0.002$). The first and the second axes explained 47.2% and 3.6%, respectively of the variability in terrestrial isopod distribution. The strongest predicting variables were number of predators (explaining 35.2%, pseudo-F = 74.8, $p = 0.002$) and the height above the ground (explaining 12.2%, pseudo-F = 31.8, $p = 0.002$), whereas time after astronomical dusk




Figure 4. Vertical distribution of terrestrial isopod species and predators on the wall per day in 90-min observation.

and air temperature explained less than 2% of the variability in the terrestrial isopod distribution (1.8%, pseudo-F = 4.8, $p = 0.002$ and 1.9%, pseudo-F = 5.3, $p = 0.01$, respectively). The effect of air humidity was not significant (0.2%, pseudo-F = 0.8, $p = 0.456$). The presence of predators is a good predictor for the presence of terrestrial isopod species (Table 1, Fig. 6A), although *P. spinicornis* was most numerous at sites with medium numbers of predators.

Based on GAM models, the number of terrestrial isopods was significantly predicted by the height of the record on the wall for all species except *P. pruinus* (Table 1), the three other species were most numerous at a height of 70–80 cm above the ground (Fig. 6B). The time after astronomical dusk was a significant predictor for the number of observed individuals of *A. versicolor* (Table 1), which reached the highest numbers ~ 3 h after dusk (Fig. 6C). The air temperature had no significant effect on the prediction of the numbers of observed individuals of single terrestrial isopod species (Table 1).

Table 1. Summary of fitted Generalised Additive Models for environmental variables predicting the numbers of observed terrestrial isopods on the wall during a 90-min observation in October 2019. Significant effects in bold. (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, n.s. not significant).

Predictors Response	predators (ind.)			height (cm)			time after sunset (min)			air temperature (°C)		
	R2[%]	F	p	R2[%]	F	p	R2[%]	F	p	R2[%]	F	p
<i>A. versicolor</i>	54.3	71.2	***	19.4	14.4	***	6.4	4.1	*	4.3	2.7	n.s.
<i>A. vulgare</i>	29.2	24.8	***	10.1	6.7	**	4.5	2.9	n.s.	4.3	2.7	n.s.
<i>P. pruinus</i>	29.3	24.8	***	3.4	2.1	n.s.	0.8	0.5	n.s.	3.1	1.9	n.s.
<i>P. spinicornis</i>	23.4	18.3	***	10.3	6.9	**	2.4	1.5	n.s.	0.7	0.4	n.s.




Figure 5. RDA-biplot for the distribution of different species of terrestrial isopods (blue arrows) on the wall during a 90-min observation in October 2019 and environmental variables (red arrows). Only environmental variables with a significant effect on terrestrial isopod distribution are presented.

Discussion

We present data on the distribution of four species of terrestrial isopods on a wall at night in autumn 2019 in the Czech Republic. The distribution of the most numerous species was significantly explained by the height on the wall and for one species there was also a significant correlation between the number of observed individuals and the time after sunset. Distribution of all species had a significant relationship with the presence of predators, which mainly consisted of spiders.

All species of terrestrial isopods found on the studied wall are common inhabitants of cities in the Czech Republic (Orsavová and Tuf 2018). The absence of *P. scaber* is surprising because it is frequently found on walls of buildings in the nearby city of Olomouc (pers. obs.) and it was reported from walls several times (e.g., Den Boer 1961), including brick walls (Meinertz 1944). The only species found in this study and reported before from (limestone) walls is *P. spiniornis* (Sutton 1972; Boeraeve et al. 2021).

The preferred height of all species was ~ 70–80 cm, with low numbers of animals at 60 cm (Fig. 4). It is necessary to say that at a height of 60 cm there was a small prominent lintel between the lower stones and the upper brick parts of the wall (see Fig. 2A). For some reason, animals (both terrestrial isopods and spiders) were more numerous above and below this brick lintel. Higher numbers of observed individuals could be related to higher humidity near the lintel (catching rainfall) or small accumulation of debris including excrements (see Fig. 2B); probably without the presence of the lintel, distribution of terrestrial isopods on the wall would be less unequal.




Figure 6. Significant response (see Table 1) of terrestrial isopod numbers per species to **A** number of predators **B** height on the wall **C** time after dusk during a 90 min interval in October 2019.

The observed temporal pattern, with the highest number of observed individuals at approximately 21:30 h, is in accordance with the temporal activity pattern observed for *P. scaber* in the Oxford study (Brereton 1957). This species was active there from 19:00 till 03:00 h, with the highest observed activity around 23 h, indicating that its activity increased continuously the first hours after sunset. Brereton (1957) observed individuals of *P. scaber* mainly on well-structured bark of oaks and sycamores. On the other hand, Den Boer (1961) reported the highest numbers of *P. scaber* after sunset and before dawn. He supposed that individuals climb the tree trunks actively each evening from litter around the tree base because they are not able to find shelter on the smooth bark of studied aspens. The wall we studied is old with partly eroded ground between the bricks with many crevices offering a lot of shelter. We do not suppose that terrestrial isopods climb on the wall from shelters at or near the soil every night and therefore our data support the view of Brereton (1957) that the terrestrial isopods find daytime shelter on the wall.

We found a weak effect of air temperature and no effect of air humidity on the activity of terrestrial isopods. Cloudsley-Thompson (1958), as well as Den Boer (1961), reported an effect of air humidity in explaining the vertical activity of terrestrial isopods. However, we did not find this effect, which may be an artefact of the low variability in air humidity during our research (58–70%, one night of 46%), but it is also necessary to say that the genus *Armadillidium* is less vulnerable to low humidity levels than, e.g., *P. scaber* (Cloudsley-Thompson 1977) and has a higher drought resistance (Dias et al. 2013). On the other hand, Brereton (1957) did not find effects of air humidity, nor air temperature, on the activity of *P. scaber*.

The strongest predictive power for the numbers of observed terrestrial isopods on the wall was the number of observed predators. Centipedes, spiders, and harvestmen are known isopod predators (Cloudsley-Thompson 1958; Sutton 1970; Santos and Gnaspini 2002). The observed predatory individuals were the centipede *Lithobius forficatus* (Linnaeus, 1758), the harvestman *Phalangium opilio* Linnaeus, 1761, and the spiders *Nuctenea umbratica* (Clerck, 1757) and *Steatoda grossa* (C. L. Koch, 1838). Of course, terrestrial isopods do not prefer places with high predatory pressure, but

the reason for the presence of predators at specific heights with high numbers of terrestrial isopods is probably due to accessibility of their prey or overall suitability of environmental conditions there. Nevertheless, we can conclude that high abundance of predators at a particular height could be a good predictor for terrestrial isopods on walls.

It is not yet clear exactly why terrestrial isopods are found on walls, but the search for algae as food source is most plausible. Brereton (1957) experimentally confirmed consumption of *Pleurococcus* algae from the tree bark by *P. scaber*. However, in the same experiment, algae were not consumed by *O. asellus* (Brereton 1957). We therefore encourage research that investigates feeding habits of terrestrial isopods on walls and on trees!

In conclusion, we observed four species of terrestrial isopods present on the wall during the several hours after sunset. They were distributed along the complete studied height (up to 2 m), but they preferred a height of ~ 0.75 m above the soil surface. Similar spatiotemporal patterns were recorded for spiders as their potential predators. We suppose that the terrestrial isopods shelter in fissures and crevices between bricks and that they are resident on the wall. It will be very useful to study their gut contents to discover what food is consumed, which will probably explain their distribution.

Acknowledgements

This study was partly supported by an internal grant of the Faculty of Science of Palacký University Olomouc (IGA_PrF_2021_014). The authors are grateful to Ondřej Machač for the identification of the spiders and to Pallieter De Smedt and one anonymous reviewer for valuable comments, suggestions, and corrections of the manuscript. Alena Dvorska kindly edited the manuscript for English.

References

- Abbott CH (1918) Reactions of land isopods to light. *Journal of Experimental Zoology* 27: 193–246. https://scholar.archive.org/work/f5utddbwmwbdllhdeutq64bsnz5y/access/ia_file/crossref-pre-1923-scholarly-works/10.1002%252Fjctb.5000412409.zip/10.1002%252Fjez.1400270204.pdf
- Alexander KNA (2008) *Platyarthrus hoffmannseggii* Brandt in arboreal ant nests (Isopoda, Oniscidea, Platyarthridae). *Bulletin of the British Myriapod and Isopod Group* 23: 15–16. <http://www.bmig.org.uk/sites/www.bmig.org.uk/files/bulletin/BullBMIG23-2008.pdf#page=17>
- Alexander KNA (2011) *Armadillidium depressum* Brandt, 1833 climbing trees in Dorset. *Bulletin of the British Myriapod and Isopod Group* 25: 37–38. <https://www.bmig.org.uk/sites/www.bmig.org.uk/files/bulletin/BullBMIG25-2011.pdf#page=39>
- Boeraeve P, Arijs G, Segers S, De Smedt P (2021) Habitat and seasonal activity patterns of the terrestrial isopods (Isopoda: Oniscidea) of Belgium. *Belgian Journal of Entomology* 116: 1–95.

- Božanič B, Hradílek Z, Machač O, Pižl V, Štáhlavský F, Tufová J, Věle A, Tuf IH (2013) Factors affecting invertebrate assemblages in bryophytes of the Litovelské luhy National Nature Reserve, Czech Republic. *Acta Zoologica Bulgarica* 65(2): 197–206.
- Brereton JLG (1957) The distribution of woodland isopods. *Oikos* 8: 85–106. <https://doi.org/10.2307/3564994>
- Cloudsley-Thompson JL (1958) The effect of wind upon the nocturnal emergence of woodlice and other terrestrial arthropods. *The Entomologist's Monthly Magazine* 94: 106–108, 184–185, 283–284.
- Cloudsley-Thompson JL (1977) *The Water and Temperature Relations of Woodlice*. Meadowfield Press Ltd., Durham, 84 pp.
- Cole LC (1946) A study of the cryptozoa of an Illinois woodland. *Ecological Monographs* 16: 49–86. <https://doi.org/10.2307/1943574>
- Den Boer PJ (1961) The ecological significance of activity patterns in the woodlouse *Porcellio scaber* Latr. (Isopoda). *Archives Néerlandaises de Zoologie* 14: 283–409. <https://doi.org/10.1163/036551661X00070>
- Dias ATC, Krab EJ, Mariën J, Zimmer M, Cornelissen JHC, Ellers J, Wardle DA, Berg MP (2013) Traits underpinning desiccation resistance explain distribution patterns of terrestrial isopods. *Oecologia* 172(3): 667–677. <https://doi.org/10.1007/s00442-012-2541-3>
- Dutton A-J, Pryce D (2018) UV fluorescence in a critically endangered isopod, *Pseudolaureola atlantica* (Vandel, 1977). *Bulletin of the British Myriapod & Isopod Group* 30: 33–38. https://www.bmig.org.uk/sites/www.bmig.org.uk/files/bulletin/Bull-BMIG30p33-38_Dutton_Pseudo-antlantica.pdf
- Fritsche H (1934) Über Wanderungen von *Porcellio scaber* (SARS). *Zoologischer Anzeiger* 107: 62–64.
- Glazier DS, Kleynhans E (2015) Arboreal herbivory by a semi-terrestrial South African isopod crustacean, *Tylos capensis* Krauss (Isopoda: Tylidae), on the Bietou Bush, *Chrysanthemoides monilifera* (L.) Norlindh. *African Invertebrates* 56: 729–738. <https://doi.org/10.5733/afin.056.0315>
- Gregory S (2009) Woodlice and Waterlice (Isopoda: Oniscidea & Asellota) in Britain and Ireland. Biological Records Centre, Wallingford, 176 pp.
- Mechthold O, Boeraeve P, De Smedt P (2021) Distribution of two terrestrial isopods (Isopoda: Oniscidea) on bark of living beech trees in an old forest reserve. In: Abstract Book of the 11th International Symposium on Terrestrial Isopod Biology. Spinicornis and ForNaLab, Department of Environment, Faculty of Bioscience Engineering, Ghent University, Belgium, 38 pp.
- Meinertz TH (1944) Beiträge zur Ökologie der Landisopoden mit besondere Berücksichtigung ihrer Atmungsorgane. *Zoologische Jahrbücher. Abteilung für Systematik, Geographie und Biologie der Tiere* 76: 501–518.
- Moeed A, Meads MJ (1983) Invertebrate fauna of four tree species in Orongorongo Valley, New Zealand, as revealed by trunk traps. *New Zealand Journal of Ecology* 6: 39–53. <https://www.jstor.org/stable/24052727>
- Orsavová J, Tuf IH (2018) Suchozemští stejnonožci: atlas jejich rozšíření v České republice a bibliografie 1840–2018 [Woodlice: Distribution Atlas in the Czech Republic and Bibliog-

- raphy 1840–2018]. Acta Carpathica Occidentalis, Supplement 1: 1–120. [in Czech with English abstract]
- Růžička V, Boháč J, Macek J (1991) Bezobratlí živočichové dutých stromů na Třeboňsku [Invertebrate animals from hollow trees in the Třeboň basin]. Sborník Jihočeského muzea v Českých Budějovicích, přírodní vědy 31: 33–46. [in Czech with English summary]
- Santos FH, Gnaspiri P (2002) Notes on the foraging behavior of the Brazilian cave harvestman *Goniosoma spelaicum* (Opiliones, Gonyleptidae). The Journal of Arachnology 30: 177–180. [https://doi.org/10.1636/0161-8202\(2002\)030\[0177:NOTFBO\]2.0.CO;2](https://doi.org/10.1636/0161-8202(2002)030[0177:NOTFBO]2.0.CO;2)
- Schmidt C (1999) Redescription of *Atracheodillo marmorivagus* (Isopoda, Oniscidea, Eubelidae), an arbicolous isopod from Congo and Rwanda. Bonner Zoologische Beitrage 48: 353–365. https://zoologicalbulletin.de/BzB_Volumes/Volume_48_3_4/353_365_BZB48_3_4_Schmidt_Christian.PDF
- Šmilauer P, Lepš J (2014) Multivariate Analysis of Ecological Data using Canoco 5. Cambridge University Press, Cambridge, 362 pp. <https://doi.org/10.1017/CBO9781139627061>
- Standen R (1921) The woodlice and other Crustacea of Derbyshire and Staffordshire. Mainly from Dovedale and District. The Lancashire and Cheshire Naturalist 13: 130–138.
- Sutton SL (1970) Predation on woodlice; an investigation using the precipitin test. Entomologia Experimentalis et Applicata 13: 279–285. <https://doi.org/10.1111/j.1570-7458.1970.tb00111.x>
- Sutton SL (1972) Invertebrate Types – Woodlice. Ginn & Co. Ltd, London, 144 pp.
- Tracz H (2013) Diplopoda, Chilopoda i Isopoda w waloryzacji ekosystemów leśnych Leśnego Kompleksu Promocyjnego “Lasy Spalsko-Rogowskie” [Diplopoda, Chilopoda and Isopoda in valorization of the forest ecosystems of the Forest Promotional Complex “Lasy Spalsko-Rogowskie”]. Studia i Materiały Centrum Edukacji Przyrodniczo-Leśnej 15(35): 36–47. [in Polish with English abstract] <https://bibliotekanauki.pl/articles/881458>
- Thunes KH, Søli GEE, Thuróczy C, Fjellberg A, Olberg S, Roth S, Coulianos C-C, Disney RHL, Starý J, Vierbergen G, Jonassen T, Anonby J, Köhler A, Menzel F, Szadziewski R, Stur E, Adaschkiewicz W, Olsen KM, Kvamme T, Endrestøl A, Podenas S, Kobro S, Hansen LO, Kvitte GM, Haenni J-P, Boumans L (2021) The arthropod fauna of oak (*Quercus* spp., Fagaceae) canopies in Norway. Diversity 13: e332. <https://doi.org/10.3390/d13070332>
- Van Name WG (1936) The American land and fresh-water isopod Crustacea. Bulletin of the American Museum of natural History 71: 1–535.
- Vandel A (1962) Isopodes terrestres (deuxième partie). Fédération Française des Sociétés de Sciences Naturelles, Paris. Faune de France 66: 417–932.
- Warburg MR (1993) Evolutionary Biology of Land Isopods. Springer-Verlag, Berlin, 160 pp. <https://doi.org/10.1007/978-3-662-21889-1>