

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/344375467>

# Extension of host range for *Anilocra dimidiata*, *Nerocila sigani* and first record of *Nerocila depressa* (Isopod: Cymothiod) from Odisha coast, India

Article in *Indian Journal of Geo-Marine Sciences* - September 2020

CITATIONS

6

READS

144

4 authors:



Jaya Kishor Seth

Berhampur university

27 PUBLICATIONS 44 CITATIONS

SEE PROFILE



Amit KUMAR Behera

Berhampur university

3 PUBLICATIONS 7 CITATIONS

SEE PROFILE



Swarup Ranjan Mohanty

Zoological Survey of India

55 PUBLICATIONS 104 CITATIONS

SEE PROFILE



Anil Mohapatra

Zoological Survey of India

176 PUBLICATIONS 973 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Meiofaunal Diversity of Coastal Midnapore(East) District,West Bengal,India [View project](#)



Studies on the population structure, and genetic diversity based on D-loop region of mitochondrial DNA, and germplasm resources mapping of near threatened and vulnerable sheatfishes (family: Siluridae) of Odisha, sponsored by DST, Govt. of Odisha [View project](#)



### Short Communication

## Extension of host range for *Anilocra dimidiata*, *Nerocila sigani* and first record of *Nerocila depressa* (Isopod: Cymothiod) from Odisha coast, India

J K Seth<sup>\*a</sup>, A K Behera<sup>a</sup>, S R Mohanty<sup>b</sup> & Anil Mohapatra<sup>b</sup>

<sup>a</sup>Post Graduate Department of Zoology, Berhampur University, Odisha – 760 007, India

<sup>b</sup>Estuarine Biology Regional Centre, Zoological Survey of India, Gopalpur-on-Sea, Ganjam, Odisha – 761 002, India

\*[E-mail: jkseth52@gmail.com]

Received 26 March 2019; revised 14 November 2019

This report of parasitic isopod *Anilocra dimidiata* from the host *Karalla daura* and *Nerocila sigani* from the host *Lutjanus lutjanus* signifies their host range extension. Further, the report of *Nerocila depressa* from the host *Selaroides leptolepis* is the first record for the state of Odisha.

[**Keywords:** Host extension, Isopod, New record, Odisha coast]

### Introduction

The isopods of the genus *Anilocra* and *Nerocila* are the external cymothiods known to inflict range of effects on the scale, pigmentation of the body surface, epidermis, bone, connective tissues and also induce behavioural changes of the host fishes<sup>1-4</sup>. The isopods of the genus *Anilocra* is comprised of 50 species, out of which three species viz., *Anilocra dimidiata* Bleeker 1857, *Anilocra longicauda* Schiödt & Meinert 1881 and *Anilocra leptosoma* Bleeker 1857 were reported from Indian water<sup>5</sup>. The species *A. dimidiata* has been reported from the host *Sardinella longiceps* Valenciennes, 1847, *Psettus* spp., *Phloptectoralis bindus* (Valenciennes, 1835), *Nemipterus* spp., *Epinephelus* spp. and *Lactarius lactarius* (Bloch & Schneider 1801)<sup>6</sup>.

The genus *Nerocila* is composed of 65 species, out of which 17 nominal species were reported from Indian water<sup>7</sup>. The isopod *Nerocila sigani* Bowman and Tareen, 1983 parasite was reported from the host fish species viz., *Siganus canaliculatus* (Park, 1797), *Parastromateus nige* (Bloch, 1795) and *Terapon theraps* Cuvier, 1829. An earlier study reported the parasitic nature of *Nerocila sigani* in the host *Terapon theraps* from the coastal water of Odisha<sup>8</sup>. The

species *Nerocila depressa* Milne Edwards, 1840 was reported from the host *Opisthopterus tardoore* (Cuvier, 1829), *Selaroides leptolepis* (Cuvier, 1833) and its geographical distribution is limited to Tamil Nadu coast<sup>7,9</sup>.

### Materials and Methods

Parasitic isopods were collected along with their host fish species (Fig. 1 A–G) during the month of December, 2018 from Haripur fish landing centre (19° 15' 43.22"N, 84° 54' 50.01" E), Gopalpur-on-Sea, Odisha coast, India. The host species *Karalla daura* (Cuvier, 1829) was parasitized by *Anilocra dimidiata*, *Lutjanus lutjanus* Bloch, 1790 and *Siganus canaliculatus* (Park, 1797) were parasitized by *Nerocila sigani* and the isopod *Nerocila depressa* was found attached to the body surface of host species *Selaroides leptolepis*. All these parasites were gently pulled out from the host body and preserved in 70 % alcohol. The parasites are deposited in the national repository of Estuarine Biology Regional Centre, Zoological Survey of India, Gopalpur-on-Sea, Odisha, India. The host fish taxonomy and nomenclature was according to FishBase<sup>10</sup>. The parasites were identified based on the standard keys and description<sup>6,7,11,12</sup>.

### Results

#### Systematics

Order: Isopoda

Family: Cymothoidae

Genus: *Anilocra*

**1. *Anilocra dimidiata*** Bleeker, 1857 (number of specimens examined: 01 female, registration number: EBRC/ZSI/Cr-10968): Total length (head to telson): 25.07 mm; total width (at 5<sup>th</sup> pereonite): 7.51 mm, number of pereonite: 07, number of pleonite: 05, length of pleotelson: 5.65 mm and width of pleotelson: 5.32 mm.

#### Description

Body is 3.33 times as elongated as width. The coxa is barely visible in dorsal view. Eyes are distinct and large with facets. Posterolateral margin is not produced at any pereonite. Sixth pereonite is longest. 1<sup>st</sup>, 5<sup>th</sup> and 7<sup>th</sup> pereonite sub-equal in length. Pleonite 1




Fig. 1 — (A-G) Cymothoid isopods with their host. A) *Anilocra dimidiata* (female) with the host *Karalla daura*. B) *Anilocra dimidiata* (female). C) *Nerocila sigani* (female) with the host *Lutjanus lutjanus*. D) *Nerocila sigani* (female) with the host *Siganus canaliculatus*. E) *Nerocila sigani* (female). F) *Nerocila depressa* with host *Selaroides leptolepis*. G) *Nerocila depressa*

is longest and pleonite 1–5 don't have posterolateral margin. Pleotelson lateral margin turned up and with caudomedial lobe. Antennule with 7 numbers of articles which are extending up to the eye. Antenna with 10 numbers of articles extending up to pereonite 2. Body is white in colour; however half of the body is covered with chromatophores.

**2. *Nerocila sigani*** Bowman & Tareen, 1983 (number of specimens examined: 03 female, registration number: EBRC/ZSI/Cr-10969): Total length (head to telson): 15.44–21.61 mm (mean: 19.54 mm); total width (at 5<sup>th</sup>–6<sup>th</sup> pereonite): 8.05–11.48 mm (mean:

9.61 mm); number of pereonite: 07; number of pleonite: 05; length of pleotelson: 4.1–5.37 mm (mean: 4.75 mm) and width of pleotelson: 3.14–5.82 mm (mean: 4.74 mm).

#### Description

Boby is 1.8–2 times as elongated as width and widest in between 5<sup>th</sup> and 6<sup>th</sup> pereonite. Cephalon is as elongated as width with rounded frontal margin. Eyes with facets almost indistinct. Pereonites 1<sup>st</sup>, 5<sup>th</sup>, 6<sup>th</sup> and 7<sup>th</sup> are subequal in length, pereonite 2–4 short and 4<sup>th</sup> is shortest. There is a depression seen on the venterolateral margin of pereonite 1–4. Coxae of pereonite 2–7 are visible in dorsal view. Pereonite coxae 6–7 are longest. Posterolateral margins produced backward. Pleonites are visible having posterolateral projections. Pleonite 1 is shortest and 2–5 are subequal in length. Pleotelson is as elongated as width smoothly rounded without caudomedial lobe. Antennule with 8 articles and antenna with 10 articles extending up to 1<sup>st</sup> pereonite. Body is white in colour and there also present a distinct venterolateral margin extending from cephalon to pleotelson.

**3. *Nerocila depressa*** Milne Edwards, 1840 (number of specimens examined: 01 female, registration number: EBRC/ZSI/Cr-10970): Total length (head to telson): 19.32 mm; total width (at 5<sup>th</sup> pereonite): 8.32 mm; number of pereonite: 07; number of pleonite: 05; length of pleotelson: 4.25 mm and width of pleotelson: 5.31 mm

#### Description

Body is about 2.3 times as elongated as width. Cephalon apex margin is rounded. Eyes are not visible. Coxae of pereonite 2–7 are visible in dorsal view. The posterolateral margins of pereonites 1–7 are produced. Fifth pereonite is the widest and longest. Pleonites are distinct with posterolateral margins produced backward. First pleotelson is shortest and 2–5 are subequal. Pleotelson is subtriangular in shape. It has two long distinct caudal projections. Antennule with 8 articles and antenna with 10 articles. Antennule and antenna extending up to the posterior portion of cephalon. Body is white in colour with Cephalon, lateral portion of the pereonites, pleonites and uropod projections are having black green chromatophores.

#### Discussion

The present study provides the first material evidence regarding the occurrence of the isopod

parasite *Anilocra dimidiata* from the host *Karalla daura* and the record of the parasite *Nerocila sigani* from the host *Lutjanus lutjanus*, which confirms the range extension of host for these parasite. Further, the report of *Nerocila depressa* from the host *Selaroides leptolepis* is the first record for the state of Odisha. These parasites are economically important as they are causing considerable damage to the fisheries resources, so it is high time to carry out a comprehensive long term study on the isopods parasites and their impact assessments on host fishes along the Odisha coast, India.

### Acknowledgements

First two authors are thankful to the authority of Berhampur University, Odisha, India for providing necessary facilities to work. Third and fourth author are thankful to the Director, Zoological Survey of India, Kolkata for providing necessary facilities for the work.

### Conflict of Interest

Authors don't have any conflict of interest.

### Author Contributions

AKB & SRM: Collection, preservation and identification and manuscript preparation. JKS & AM: Identification, manuscript preparation and critical analysis.

### References

- 1 Bowman T E & Mariscal R N, *Nerocila heterozota*, a new cymothoid isopod, with notes on its host, the anemone fish, *Amphiprion akallopisos*, in the Seychelles, *Crustaceana*, 14 (1968) 97-104.
- 2 Morton B, Host specificity and position on the host in *Nerocila phaeopleura* Bleeker (Isopoda, Cymothoidae), *Crustaceana*, 26 (1974) 143-148.
- 3 Williams L B & Williams E H, Nine new species of *Anilocra* (Crustacea: Isopoda: Cymothoidae) external parasites of West Indian coral reef fishes, *P Biol Soc Wash*, 94 (1981) 1005-1047.
- 4 Adlard R D & Lester R J G, Dynamics of the interaction between the parasitic isopod, *Anilocra pomacentri*, and the coral reef fish, *Chromis nitida*, *Parasitology*, 109 (1994) 311-324.
- 5 Aneesh P T, Helna A K, Trilles J P & Chandra K, Occurrence and redescription of *Anilocra leptosome* Bleeker, 1857 (Crustacea: isopoda: Cymothoidae) parasitizing the clupeid fish *Tenuelosatoli* (Valenciennes) from the Arabian Sea, India. *Mar Biodivers*, 49 (1) (2019), 443-450. doi: 10.1007/s12526-017-0828-7.
- 6 Rameshkumar G, Ravichandran S & Trilles J P, Cymothoidae (Crustacea, Isopoda) from Indian fishes, *Acta Parasitol*, 56 (2011) 78-91.
- 7 Trilles J P, Rameshkumar G & Ravichandran S, *Nerocila* species (Crustacea, Isopoda, Cymothoidae) from Indian marine fishes, *Parasitol Res*, 112 (2013) 1273-1286.
- 8 Dev Roy M K & Mitra S, New host record for *Nerocila sigani* (Isopoda: Cymothoidae) from Odisha coast, India, *Curr Sci*, 104 (2013) 1134-1135.
- 9 Rameshkumar G & Ravichandra S, Histopathological changes in the skins and gills of some marine fishes due to parasitic isopod infestation, *J Coast Life Med*, 1 (2013) 12-18.
- 10 Froese R. & Pauly D (eds.) *FishBase*, World Wide Web electronic publication. [http:// www.fishbase.org](http://www.fishbase.org) Version (06/2018).
- 11 Bruce N L, Australian species of *Nerocila* Leach, 1818, and *Creniola* n. gen., (Isopoda: Cymothoidae), Crustacean parasites of marine fish, *Rec Aust Mus*, 39 (1987) 355-412.
- 12 Bruce N L & Harrison-Nelson E B, New Records of fish parasitic Marine isopod crustaceans (Cymothoidae, Subfamily Anilocrinae) from the Indo-West Pacific, *P Biol Soc Wash*, 101 (1998) 585-602.