

Genetic diversity of bacteria associated with the hindgut of the terrestrial crustacean *Porcellio scaber* (Crustacea: Isopoda)

Rok Kostanjšek^a, Jasna Štrus^a, Gorazd Avguštin^{b,*}

^a University of Ljubljana, Biotechnical Faculty, Biological Department, Večna pot 111, 1111 Ljubljana, Slovenia

^b University of Ljubljana, Biotechnical Faculty, Zootechnical Department, Groblje 3, 1230 Domžale, Slovenia

Received 20 June 2001; received in revised form 3 February 2002; accepted 6 February 2002

First published online 9 April 2002

Abstract

Molecular approaches were used to examine the genetic diversity of bacteria associated with the gut wall of the terrestrial isopod *Porcellio scaber* and to determine whether an autochthonous microflora exists in the *P. scaber* hindgut. 16S ribosomal genes were amplified from the total DNA isolated from thoroughly washed papillate regions of the hindgut, where the highest concentrations of bacteria are commonly found. The amplified genes were cloned, sequenced and phylogenetically analysed. The results implied an unexpectedly large diversity of microflora associated with the cuticle of the hindgut. Almost half of the retrieved sequences were found to be less than 80% homologous with any of the known sequences available at DNA data banks. Most of these sequences were clustered in one of three groups, and were clearly distant from the sequences of other bacterial taxa, indicating that they could represent novel bacterial species or even genera. More than two thirds of the sequences were found to be phylogenetically related to sequences from bacteria typically isolated from human and animal intestines, e.g. streptococci, enterococci, and members of the genus *Bacteroides*. The majority of the remaining sequences were most closely related to typical soil bacteria, e.g. bacilli and pseudomonads. The facts that a large proportion of the retrieved sequences was related to the sequences of bacteria, which are autochthonous to intestinal ecosystems, and that bacteria, specifically attached to the cuticular spines, were observed, indicate that truly autochthonous bacteria may well be present in the hindgut of *P. scaber*. © 2002 Published by Elsevier Science B.V. on behalf of the Federation of European Microbiological Societies.

Keywords: Terrestrial isopod; Hindgut; Intestinal bacteria; Indigenous microbial flora; Genetic diversity; 16S rDNA

1. Introduction

The common woodlouse *Porcellio scaber*, Latreille 1804 (Crustacea: Isopoda) is a widely distributed terrestrial isopod. Terrestrial isopods are believed to play an important role in decomposition processes and global carbon cycling. In spite of its omnivorous nutritional habits, the diet of *P. scaber*, as that of most terrestrial isopods, consists mainly of decayed plant material composed of cellulose and other polysaccharides [1]. Although the role of microorganisms is clearly crucial in the process of degradation of complex polysaccharides such as cellulose, it has not been as thoroughly investigated in the intestinal tract of *P. scaber* as in some other arthropods which feed on cellulose-rich diets [2,3].

The digestive system of *P. scaber* consists of a foregut (stomodeum) and a hindgut (proctodeum), both lined with a cuticle. The midgut (mesenteron) is reduced to two pairs of tubular midgut glandular caeca, also known as hepatopancreas, which is the key site for the secretion of digestive enzymes and for the absorption of soluble nutrients [4] (Fig. 1). Fragments of food pass through the foregut where they are briefly masticated, compressed and filtered into the hindgut. During this process, the food is mixed with secretions derived from the hepatopancreas, before passing into the long hindgut. In contrast to other arthropods, the hindgut of isopods is a simple straight tube subdivided along its length into an anterior region, with a pair of dorsal channels (a typhlosole), a median papillate region, which ends with a muscular sphincter, and a short posterior rectum. Absorptive properties have been attributed to both the anterior hindgut and the hepatopancreas [5–7] whereas in the papillate region water and ions are reabsorbed [8–10].

Microbes in the gut of terrestrial isopods have been

* Corresponding author. Tel.: +386 (1) 721 78 27;

Fax: +386 (1) 724 10 05.

E-mail address: gorazd.avgustin@bfro.uni-lj.si (G. Avguštin).




Fig. 1. Anatomy of the digestive system of an adult *P. scaber*. ST – stomach, MG – midgut glands (hepatopancreas), ANT – anterior region of the hindgut, TI – typhlosole, PAP – papillate region of the hindgut, SP – sphincter, RE – rectum, AN – anus. Scale bar: 5 mm.

observed and studied by several authors [11–20]. Their presence indicates that the isopod gut represents a suitable environment for microbial colonisation, although it lacks specialised structures or modifications of the gut, which would facilitate colonisation [11,12,14]. The majority of the microbial flora is found in the hindgut, whereas the foreguts of terrestrial isopods are generally poorly colonised [15]. Data on the microflora inhabiting the digestive system of terrestrial isopods were acquired mainly through observations by means of light and electron microscopy, and by isolation and further chemotaxonomic studies of microorganisms from food, intestines and faeces of the animal [11–14,16–20]. Culture-dependent methods are biased, however, since the proportion of cultivable microorganisms in natural environments is often very low [21]. Such approaches are also not suitable if one wants to determine whether a part of the microbial community is truly autochthonous to the given environment.

Data concerning the presence of autochthonous bacterial flora in the isopod gut are contradictory. According to Dubos et al. [22] the term autochthonous flora is appropriate for describing the microorganisms resident in the gut and possessing mechanisms which enable them to adapt to the gut environment. These mechanisms were developed during co-evolution with the host, and do not occur in other environments. Autochthonous microorgan-

isms can live either in a mutualistic or commensalistic relationship with the host [23]. Different authors have used various terms to describe microorganisms discovered in the isopod gut. Whereas some authors support the idea of the presence of autochthonous microbial flora in the isopod gut [12–14], others suggest the absence of such flora [6,15] as a consequence of the simple anatomy of the gut, the short retention time of the food and the moulting of the hindgut cuticle, which occurs approximately once a month.

In the work reported here, cloning and sequencing of PCR-amplified 16S rRNA genes was applied in order to screen the genetic diversity of bacteria resident in the *P. scaber* hindgut and to resolve their taxonomic status. It was anticipated that some of the bacteria associated with the cuticular lining of the papillate region, where the highest density of microorganisms occurs, could represent the autochthonous bacterial flora, which has not yet been proven.

2. Materials and methods

2.1. Animals

The isopods examined in this study were collected in the spring of 1999 in woods near Cerknica in Slovenia. The animals were kept at 20°C in glass tanks filled with soil, under conditions of high humidity and 12–12 h day/night illumination. The animals were fed on leaf litter from the collection site for at least 1 month prior to the experiment. Adult animals of both sexes were used in the experiment.

2.2. Gut preparation and DNA extraction

The guts of five healthy adult isopods were extracted with fine-tipped sterile forceps and the papillate regions of the hindguts were removed. The papillate regions were opened with a sterile needle, transferred to a 0.5-ml sterile phosphate-buffered saline (PBS buffer; 130 mM NaCl, 3 mM NaH₂PO₄, 7 mM Na₂HPO₄; [pH 7.4]), and washed vigorously for 10 s. Washing was repeated three times.

Total DNA was extracted from the isolated papillate regions by the modified method proposed by Mäntynen and Lindström [24]. The papillate regions and the attached microorganisms were crushed using a teflon homogeniser in 0.5 ml of PBS buffer. Tissue particles and bacterial cells were harvested by centrifugation and then resuspended in 100 µl of TE buffer (10 mM Tris–HCl [pH 8.0], 1 mM EDTA) in the presence of 1 µl of proteinase K (20 mg ml⁻¹) and 8 µl of lysozyme (50 mg ml⁻¹). Following incubation at 37°C for 1 h, 60 µl of 10% sodium dodecyl sulfate and 20 µl of 10% *N*-lauroylsarcosine were added, and the mixture was incubated for further 30 min at 37°C. A 60 µl aliquot of preheated (65°C) 10% CTAB in 0.7%

NaCl was then added, and the mixture was incubated for 10 min at 65°C. The sample was further treated by adding an equal volume of phenol–chloroform–isoamyl alcohol (25:24:1), and then centrifuged at 10 000 × *g* for 20 min. The DNA was precipitated with an equal volume of isopropyl alcohol, centrifuged at 20 000 × *g* for 15 min, and rinsed with 70% ethanol. After centrifuging at 13 000 × *g* for 10 min, the DNA was air dried and resuspended in 20 µl of the TE buffer, pH 8.0.

2.3. PCR amplification and cloning of 16S rDNA

16S rRNA genes from total DNA were amplified using the modified evolutionary conserved eubacterial primers fD1 [25] and 1392r [26] as described elsewhere [27] with minor modifications. The sample DNA (1 µl) was amplified in a reaction mixture that consisted of 1 × reaction buffer (Gibco BRL, Carlsbad, CA, USA), 2.5 mM MgCl₂, 0.2 mM of each dNTP, 6 pmol of each primer and 1 U of *Taq* polymerase (Gibco BRL) in a final volume of 20 µl. The reaction mixture was first denatured at 94°C for 5 min, and then subjected to 30 PCR cycles (40 s at 94°C, 30 s at 60°C, and 80 s at 72°C) followed by 10 min at 72°C.

The amplified 16S rRNA genes were separated by agarose gel electrophoresis, cut from the gel and purified using a QIAquick Gel Extraction Kit (Qiagen, Germantown, MD, USA), according to the manufacturer's recommendations. The purified amplicons were digested with *Sal*I (Promega, Madison, WI, USA) and *Not*I (New England Biolabs, Beverly, MA, USA) restriction endonucleases, as recommended by the manufacturers, and subsequently cloned in the phagemid pBluescript KS II+ (Stratagene, La Jolla, CA, USA). The ligated vectors carrying 16S rRNA amplicons were used for the transformation of *Escherichia coli* DH-5-α recipient cells (Gibco BRL) by the CaCl₂ method [28].

2.4. Restriction fragment length polymorphism (RFLP) analysis and sequencing

The white colonies grown on LB/xgal plates were further analysed by restriction analysis. The recombinant plasmids were purified by a 'mini-prep method' with 5% CTAB and ethanol precipitation [28], and digested with *Taq*I or *Dde*I (Boehringer Mannheim, Germany) restriction endonucleases according to the manufacturer's recommendations. Jaccard's coefficients were calculated on the basis of the observed RFLP profiles from 29 randomly selected recombinant plasmids, and a phenogram was constructed with the UPGMA method, using the NTSYS pc 1.80 package (Applied Biostatistics Inc.) [29]. Representative clones from established groups were chosen for further analysis, and the cloned inserts were sequenced with the fD1 sequencing primer [25] by Microsynth GmbH (<http://www.microsynth.ch/>) or Colorado State University, Department of Biochemistry (<http://mmr.bmb.colostate.edu/dna/dnaseq.html>) on our request. Some of the cloned inserts were also sequenced with the sequencing primers 338f [21] and 1100r [26], thus providing almost total sequence length information.

2.5. Phylogenetic analysis and nucleotide sequence accession numbers

The phylogenetic analysis was done essentially as described previously [27], using programmes available at the Ribosomal Database Project site (<http://www.cme.msu.edu/RDP/html/analyses.html>) [30], BLAST and FASTA homology search algorithms [31,32], and the CLUSTAL X [33] and PHYLIP (Phylogeny Inference Package version 3.57c) [34] packages. The nucleotide sequence data have been deposited in the GenBank nucleotide sequence database under accession numbers AF395310–AF395327.

3. Results

Following the amplification and cloning of the 16S rRNA genes, 29 randomly selected clones, carrying recombinant plasmids with inserts of the expected size, i.e. 1400 bp, were subjected to RFLP analysis. The RFLP analysis showed that most clones clustered in one of three larger groups (not shown). The first group (RFLP-1) consisted of 10 clones, whereas the second and the third group (RFLP-2 and RFLP-3) consisted of six clones each. The rest of the clones remained ungrouped, indicating the large diversity of the amplified and cloned 16S rRNA genes and thus of the bacteria associated with the gut wall of *P. scaber*.

A total of 18 clones representing different groups and lineages in the RFLP dendrogram were chosen for partial sequencing with the fD1 sequencing primer. The names of the obtained sequences were created in a way that various alphabetical letters were attached to the common prefix RPKs. Sequences with the highest homologies to cloned sequences were retrieved from DNA data bank using homology search algorithms. The sequences were then pairwise aligned, and similarity values were calculated (Table 1). Almost half of the analysed sequences (8 out of 19) were less than 80% similar to the most homologous sequence deposited in the data bank, whereas only three sequences were more than 95% similar to the best-matched sequence from the data bank.

The obtained sequences were then further compared with sequences of bacteria from known taxa, and a phylogenetic tree was constructed (Fig. 2). The phylogenetic analysis clustered most of the cloned 16S rRNA sequences in similar groups to those of the RFLP analysis. The sequences of eight clones from the group RFLP-1 (RKP_sBD, RKP_sAL, RKP_sB, RKP_sA, RKP_sAV, RKP_sG, RKP_sAA and RKP_sAN) formed a deeply branched cluster (Fig. 2)


Table 1

List of the nearest known sequences to the hindgut clones (named RKPs plus additional letters) obtained from the data bases

Clone	Group	Closest known sequences	Percentage of similarity	Length of aligned sequences
RKPpA	RFLP-1	<i>Macrococcus caseolyticus</i> (Y15711)	74.9	928
		<i>Staphylococcus caseolyticus</i> (D83359)	74.9	925
		<i>Streptococcus hyointestinalis</i> (AF201898)	74.8	928
RKPpB	RFLP-1	U.C. bacterium (firmicute) (AF141540)	80.9	513
		<i>Streptococcus waiu</i> (AF088900)	77.1	513
		<i>Streptococcus hyointestinalis</i> (AF201898)	76.8	514
RKPpG	RFLP-1	<i>Macrococcus carouzelicus</i> (Y15713)	72.7	821
		<i>Macrococcus equipercius</i> (Y15712)	72.6	821
		<i>Staphylococcus caseolyticus</i> (D83359)	72.6	821
RKPpAA	RFLP-1	<i>Staphylococcus sciuri</i> (S83569)	78.1	657
		<i>Macrococcus bovicus</i> (Y15714)	77.5	657
		<i>Macrococcus carouzelicus</i> (Y15713)	77.4	657
RKPpAL	RFLP-1	U.I. bacterium (soil) (AJ011421)	77.9	938
		<i>Enterococcus hirae</i> (AF061011)	74.9	938
		<i>E. faecium</i> (AF070223)	74.8	938
RKPpAN	RFLP-1	<i>Bacillus benzovorans</i> (Y14693)	73.1	940
		<i>Bacillus psychrodurans</i> (AJ277984)	73.5	940
		U.I. bacterium (hizosphere) (AJ011421)	77.5	687
RKPpAV	RFLP-1	<i>S. equinus</i> (AB002514)	78.1	739
		<i>Streptococcus hyointestinalis</i> (AF201898)	77.6	739
		<i>Streptococcus waiu</i> (AF088900)	74.7	822
RKPpBD	RFLP-1	<i>Bacillus psychrodurans</i> (AJ277984)	79.8	584
		<i>Bacillus psychrotolerans</i> (AJ277983)	79.5	577
		<i>Bacillus benzovorans</i> (Y14693)	79.2	584
RKPpI	RFLP-2	<i>B. fusiformis</i> (M77486)	77.0	820
		U.C. bacterium (manure pit) (AF261818)	84.2	495
		U.I. bacterium (soil) (AF010085)	84.3	461
RKPpS	RFLP-2	<i>B. fusiformis</i> (M77486)	76.0	831
		U.C. bacterium (manure pit) (AF261818)	86.1	489
		U.I. bacterium (soil) (AF010085)	85.4	457
RKPpZ	RFLP-2	U.I. bacterium (manure pit) (AF261818)	92.3	505
		U.I. bacterium (soil) (AJ232784)	92.2	510
		<i>B. psychrophilus</i> (D16277)	92.0	501
RKPpN	RFLP-3	<i>P. fluorescens</i> (AF094725)	83.1	958
		U.I. bacterium (sea sediment) γ Proteobacteria (AB015575)	83.1	958
		<i>Pseudomonas putida</i> (D86001)	83.0	958
RKPpAC	RFLP-3	<i>P. fluorescens</i> (AF094725)	94.0	769
		U.I. bacterium (sea sediment) γ Proteobacteria (AB015575)	94.0	769
		<i>Pseudomonas putida</i> (D86001)	93.7	763
RKPpAG	<i>Enterococcus</i>	<i>E. faecium</i> (AJ276355)	99.1	1415
		<i>Enterococcus durans</i> (AJ276354)	98.9	1418
		<i>Enterococcus hirae</i> (AJ276356)	98.7	1417
RKPpBA	<i>Enterococcus</i>	<i>E. saccharolyticus</i> (Y18357)	96.8	1695
		<i>Enterococcus gallinarum</i> (AF039898)	96.3	1395
		Eubacterium sp. (AF135452)	96.2	1405
RKPpAM	Bacteroides	U.I. bacterium (river organic aggregate) (AF150715)	93.6	1033
		U.I. bacterium (AF227834)	88.4	1418
		<i>Bacteroides acidofaciens</i> (AB021159)	87.9	1419
RKPpF	<i>Neisseria</i>	<i>Neisseria perflava</i> (AJ239295)	97.7	478
		<i>Neisseria mucosa</i> (AJ239279)	97.7	476
		<i>Neisseria flavescens</i> (L06168)	96.1	512
RKPpT		U.I. bacterium (anaerobic consortia) (AJ249104)	81.9	482
		<i>C. thermocellum</i> (L09173)	80.1	482
		<i>Acetovibrio cellulolyticus</i> (L35516)	79.6	482

The percentage of similarity was calculated on the basis of pairwise alignment. U.C., uncultivated; U.I., unidentified.

Fig. 2. Phylogenetic tree including partial 16S rDNA sequences retrieved from hindgut clones (named RKPs plus additional letters) and the closest sequences to hindgut clones from the clone library. The 16S rDNA sequence of *Aquifex pyrophilus* was used as the outgroup. The scale bar represents 0.1 substitution per nucleotide position. The numbers at the nodes indicate bootstrap values based on 1000 resamplings (values below 60 are not shown).






Fig. 3. Phylogenetic tree showing the relationships among 16S rDNA sequences from two hindgut clones, RKP sAG and RKP sBA, and the members of the genus *Enterococcus*. The 16S rDNA sequence of *E. coli* was used as the outgroup. The scale bar represents 0.1 substitution per nucleotide position. The numbers at the nodes indicate bootstrap values based on 1000 resamplings (values below 60 are not shown).

with the nearest known taxon being the genus *Streptococcus*. The similarity values to the best-matched sequences from the data bank, obtained by the pairwise sequence alignment analysis, were all below 80%. The unique line of descent was further strengthened by fairly high confidence values as revealed by the bootstrap analysis.

Three clones from the group RFLP-2 (RKP sS, RKP sI and RKP sZ) were grouped together again, as with the RFLP analysis, and the phylogenetically closest sequences were from unidentified bacteria isolated from the manure pit (AF261818) and soil (AJ232784) [35]. Sequences from this group were clustered together with sequences from members of the genus *Bacillus*, which was again supported by relatively high (91%) bootstrap values.

Two sequences from the RFLP-3 group (RKP sAC and RKP sN) clustered together with sequences from the bacteria of the genus *Pseudomonas*. The bootstrap value of 100% for the node clustering clones RKP sAC, RKP sN and members of the genus *Pseudomonas*, supports the assignment of these clones to the *Pseudomonas* group (Fig. 2).

The sequences from the clones RKP sAG, RKP sBA and RKP sAM were, as in the case of group RFLP-1, related to bacterial genera commonly inhabiting the gut of vertebrates and insects rather than to typical soil bacteria. Sequences from the clones RKP sAG and RKP sBA were placed within the cluster comprising sequences from the genus *Enterococcus*, whereas the sequence from the clone RKP sAM was placed within the cluster of sequences from the genus *Bacteroides* (Fig. 2). Since some members of the genus *Enterococcus* and most of the bacteria from the genus *Bacteroides* are commonly found in higher animals as part of the autochthonous intestinal flora, the cloned 16S rDNAs were further sequenced with additional sequencing primers. The more accurate phylogenetic positions of these three clones were therefore determined on the basis of almost complete sequences of their 16S RNA genes.

The sequence from the clone RKP sAG shared the highest homology with the sequence from *Enterococcus faecium* (99.1%), whereas the RKP sBA insert shared the highest homology with the sequence from *Enterococcus saccharolyticus* (96.8%) (Table 1). This was further confirmed by the phylogenetic analysis (Figs. 2 and 3), and supported by high confidence values (86 and 100% respectively).

The ribosomal gene sequence from the clone RKP sAM was most similar to the sequence from an unidentified bacterium from Elbe river organic aggregate (AF150715) [36]. These two sequences were placed within the *Cytophaga–Flexibacter–Bacteroides* phylogenetic group (Fig. 4), and they clearly belong to the *Bacteroides fragilis* subgroup as defined by the Ribosomal Database Project characterisation [30]. Within the *B. fragilis* subgroup, the RKP sAM and the Elbe river isolate formed a deep and distinct lineage supported by a high confidence level (85%).

Low sequence similarity (below 82%) between the sequence of the clone RKP sT and the best-matched sequences from the data bank (Table 1) suggests that this organism is very distant to major known bacterial taxa. Deep branching of the sequence (Fig. 2) indicates that RKP sT, same as members of the RFLP-1 group, could represent a distinct and evolutionary divergent taxonomic group within the domain Bacteria.

4. Discussion

The hindgut of various soil invertebrates is a favourable environment for bacterial attachment, activation and growth [37,38]. These animals are considered to form 'hot-spots' of enhanced bacterial growth, activity and species interactions in the soil habitat [37] and are therefore very important partners in the efficient decomposition of organic matter and in mineral cycling. One example, the common woodlouse *P. scaber*, has also often been as-




Fig. 4. Phylogenetic tree showing the relationships between 16S rDNA sequence from the hindgut clone RKP SAM and the members of the *Cytophaga*–*Flexibacter*–*Bacteroides* group. The 16S rDNA sequence of *E. coli* was used as the outgroup. The scale bar represents 0.1 substitution per nucleotide position. The numbers at the nodes indicate bootstrap values based on 1000 resamplings (values below 60 are not shown).

sumed to play an important role in the fragmentation of litter and in the incorporation of decomposition products in the soil profile.

Since it is assumed that terrestrial isopods do not possess endogenous cellulases, these crucial depolymerising enzymes must originate from microorganisms that are either indigenous to the ecosystem or ingested with food. The majority of authors support the latter [6,17] and some have even suggested that the gut of the woodlouse is actually not a favourable environment for microbial growth, as demonstrated by studies on the typical soil bacterium *Pseudomonas fluorescens* [19,20,39,40]. However, a process of intensified bacterial growth was observed in the faeces of *P. scaber*, and this is assumed to be the net effect of lysis during food transit through the gut and growth in the faeces after voiding. This process appears to be initiated within the hindgut [41]. Also, the retention of ingested food particles for prolonged periods of time in the hindgut of *P. scaber* is probable, as fluorescent microbeads were found in the faeces up to 16 days after feeding [20], indicating that there is enough time for

microbial growth to occur. The beads were caught in the structure of the gut, between the folds of the exocuticle lining the hindgut region and trapped in mucopolysaccharides within the hindgut. It is therefore likely that some bacterial cells become trapped in the hindgut structures too, and develop into microcolonies in such niches. Other bacterial cells may have developed specific associations with the gut wall. Drobne [16] showed that the majority of bacteria associated with the gut wall of *P. scaber* are found in the papillate region of the hindgut. Bacterial cells remained associated with the gut wall even after several consecutive and vigorous washings and some of them were attached specifically to the cuticular spines protruding into the lumen of the gut. The existence of autochthonous microflora is, in the light of the observed attachment, therefore quite plausible.

The aim of the present study was to examine the taxonomic structure of the microbial community inhabiting the gut of *P. scaber* by molecular means. The investigation was focused on microorganisms that are potentially associated with the gut wall and are capable of remaining in the gut after the removal of the gut contents and thorough washing of the inner surface. To our knowledge, a molecular investigation of the structure of the microbial community inhabiting the *P. scaber* gut has not yet been employed. By such means autochthonous bacteria, or novel bacterial species or even genera, living in the soil and proliferating in the gut of the isopods could be discovered. The latter possibility actually seemed very probable due to the known low culturability of soil bacteria [42,21].

Phylogenetic analysis indicated the presence of a diverse bacterial flora in the *P. scaber* hindgut. The RFLP analysis was applied first, and it was found that most of the cloned sequences can be clustered in three groups, whereas approximately 25% remained ungrouped. This finding was strongly confirmed by sequence analysis of the representative clones, indicating the suitability of the RFLP approach in the initial steps of such studies. Almost half of the amplified rRNA genes were less than 80% homologous with the best-matched sequences deposited in the DNA data bank, indicating that they come from bacteria that have not yet been cultivated and very likely represent novel bacterial species or even genera. This is in agreement with observations of low percentage of cultivable bacteria from a variety of natural microbial ecosystems, such as soil [21,42,43].

More than 70% of the analysed rRNA genes were found to be homologous with the ribosomal genes from Gram-positive bacteria. Eight out of 18 rRNA genes positioned together and formed a deeply branched cluster, distantly related to the genus *Streptococcus*. Some of the most closely related streptococci, e.g. *Streptococcus bovis* and *Streptococcus equinus* are anaerobes that are typically isolated from the intestines of man and animals [44–46]. The same is true for the enterococci, that were most closely related to two of the analysed clones (RKP SAM and

RKPsBA), and for the members of the genus *Bacteroides*, that were most closely related to the rRNA gene sequence RKPsAM. Hence, 11 out of 18 retrieved rRNA gene sequences were related to bacteria that typically inhabit the animal gastrointestinal tract and are generally unable to survive outside their animal host. This could indicate that a specific, presumably anaerobic microflora exists in the hindgut of *P. scaber*, and that a constant, i.e. symbiotic, type of interaction between these microorganisms and the host has developed.

Most of the remaining recovered rDNA sequences were most closely related to typical soil bacteria. Three out of six ribotypes were sequenced from the RFLP-2 group, and their closest relatives were uncultivated bacteria from soil or the manure pit (Table 1). The cultivated bacteria with the most homologous 16S rDNA sequences to those of the RFLP-2 group were *Bacillus fusiformis* and *Bacillus psychrophilus*. However, the retrieved ribotypes from *P. scaber* formed a distinct group in the phylogenetic tree together with rDNA sequences from other uncultivable bacteria distantly related to cultivable bacilli. Two out of six ribotypes were sequenced from the RFLP-3 group and they were related to rDNA sequences from bacteria belonging to the genus *Pseudomonas*. The two remaining sequences, i.e. RKPsT and RKPsF, were most homologous with the rDNA sequences from uncultivated anaerobic bacteria and *Clostridium thermocellum*, and to species from the genus *Neisseria*, respectively. The former sequence, in particular, appears to be important due to the known cellulolytic potential of the anaerobic thermophilic clostridia.

Different bacteria resisted the applied washing procedure and we may speculate that they could also withstand the rapid flow of the food through the hindgut or the moulting of the hindgut cuticle. We can therefore assume that these bacteria may be capable of remaining in the gut of *P. scaber* for prolonged periods of time. Since a good proportion of the recovered ribotypes are related to bacteria that typically inhabit the human or animal gut and represent the true autochthonous microflora of these ecosystems, the existence of autochthonous bacteria within the gut of *P. scaber* is plausible. The contradictory evidence has been acquired with traditional microbiological approaches and is likely biased due to the dependence of these methods on in vitro cultivation. It appears possible that the hindgut of *P. scaber* is inhabited by truly autochthonous bacteria that are fastidious and do not grow under defined laboratory conditions.

Acknowledgements

This work was supported by grants from the Ministry of Education, Science and Sport of the Republic of Slovenia.

References

- [1] Wieser, W. (1966) Cooper and the role of isopods in degradation of organic matter. *Science* 153, 67–69.
- [2] Breznak, J.A. and Brune, A. (1994) Role of microorganisms in the digestion of lignocellulose by termites. *Annu. Rev. Entomol.* 39, 453–487.
- [3] Cruden, D.L. and Markovetz, A.J. (1987) Microbial aspects of the cockroach hindgut. *Arch. Microbiol.* 138, 131–139.
- [4] Holdich, D.M. and Ratcliffe, N.A. (1970) A light and electron microscope study of the hindgut of the herbivorous isopod *Dynamene bidentata*. *Z. Zellforsch.* 111, 209–227.
- [5] Hames, C.A.C. and Hopkin, S.P. (1989) The structure and function of the digestive system of terrestrial isopods. *Zool. Lond.* 217, 599–627.
- [6] Hassal, M. and Jennings, J.B. (1975) Adaptive features of gut structure and digestive physiology in the terrestrial isopod *Philoscia mucosorum* (Scopoli) 1763. *Biol. Bull.* 149, 348–364.
- [7] Hryniewiecka-Szyfter, Z. and Storch, V. (1986) The influence of starvation and different diets on the hindgut of isopoda (*Mesidotea entomon*, *Oniscus asellus*, *Porcellio scaber*). *Protoplasma* 134, 53–59.
- [8] Vernon, G.M., Herold, L. and Witkus, E.R. (1974) Fine Structure of the Digestive Tract Epithelium in the Terrestrial Isopod, *Armadillidium vulgare*. *J. Morphol.* 144, 337–360.
- [9] Palackal, T., Faso, L., Zung, J.L., Vernon, G. and Witkus, R. (1984) The ultrastructure of the hindgut epithelium of terrestrial isopods and its role in osmoregulation. *Symp. Zool. Soc. Lond.* 53, 185–198.
- [10] Storch, V. and Štrus, J. (1989) Microscopic anatomy and ultrastructure of the alimentary canal in terrestrial Isopods. *Monit. Zool. Ital.* Monogr. 4, 105–126.
- [11] Reyes, V.G. and Tiedje, J.M. (1976) Ecology of the gut microbiota of *Tracheoniscus rathkei* (Crustacea, Isopoda). *Pedobiologia* 16, 67–74.
- [12] Ineson, P. and Anderson, J.M. (1985) Aerobically isolated bacteria associated with the gut and faeces of the litter feeding macroarthropods *Oniscus asellus* and *Glomeris marginata*. *Soil Biol. Biochem.* 17, 843–849.
- [13] Gunnarson, T. and Tunlid, A. (1986) Recycling of faecal pellets in isopods: microorganisms and nitrogen compounds as potential food for *Oniscus asellus* L.. *Soil Biol. Biochem.* 18, 595–600.
- [14] Griffiths, B.S. and Wood, S. (1985) Microorganisms associated with the hindgut of *Oniscus asellus* (Crustacea, Isopoda). *Pedobiologia* 28, 377–381.
- [15] Bignell, D.E. (1984) The arthropod gut as an environment for microorganisms. In: *Invertebrate–Microbial Interactions* (Anderson, J.M., Ed.), pp. 205–227. Cambridge University Press, Cambridge.
- [16] Drobne, D. (1995) Bacteria adherent to the hindgut of terrestrial isopods. *Acta Microbiol. Immunol. Hung.* 42, 45–52.
- [17] Kukor, J.J. and Martin, M.M. (1986) The effect of acquired microbial enzymes on assimilation efficiency in the common woodlouse *Tracheoniscus rathkei*. *Oecologia* 69, 360–366.
- [18] Gunnarson, T., Sundin, P. and Tunlid, A. (1988) Importance of leaf litter fragmentation for bacterial growth. *Oikos* 52, 303–308.
- [19] Ullrich, B., Storch, V. and Schairer, H. (1991) Bacteria in the food, in the intestine and on the faeces of the woodlouse *Oniscus asellus* (Crustacea, Isopoda). *Pedobiologia* 35, 41–51.
- [20] Clegg, C.D., Anderson, J.M. and Lappin-Scott, H.M. (1996) Biophysical processes affecting the transit of a genetically-modified *Pseudomonas fluorescens* through the gut of the woodlouse *Porcellio scaber*. *Soil Biol. Biochem.* 28, 997–1004.
- [21] Amann, R.L., Ludwig, W. and Schleifer, K.-H. (1995) Phylogenetic identification and in situ detection of individual microbial cells without cultivation. *Microb. Rev.* 59, 143–169.
- [22] Dubos, R., Schaedler, R.W., Costello, R. and Hoet, P. (1965) Indigenous, normal and autochthonous flora of the gastrointestinal tract. *J. Exp. Med.* 122, 67–75.

- [23] Dodds, W.K. and Henebry, G.M. (1996) The effect of density dependence on community structure. *Ecol. Model.* 93, 33–42.
- [24] Mäntynen, V. and Lindström, K. (1998) A rapid PCR-base DNA test for enterotoxic *Bacillus cereus*. *Appl. Environ. Microbiol.* 64, 1634–1639.
- [25] Weisburg, W.G., Barms, S.M., Pelletier, D.A. and Lane, D.J. (1991) 16S ribosomal DNA amplification for phylogenetic study. *J. Bacteriol.* 173, 697–703.
- [26] Lane, D.J. (1991) 16S, 23S rRNA sequencing. In: *Nucleic Acid Techniques in Bacterial Systematics* (Stackebrandt, E., Goodfellow, M. Eds.), pp. 115–148. John Wiley and Sons, Chichester.
- [27] Ramšak, A., Peterka, M., Tajima, K., Martin, C.J., Wood, J., Johnson, M.E.A., Aminov, R.I., Flint, H.J. and Avguštin, G. (2000) Unravelling the genetic diversity of ruminal bacteria belonging to the CFB phylum. *FEMS Microbiol. Ecol.* 33, 69–79.
- [28] Sambrook, J., Fritsch, E.F., Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual*, 2nd edn. Cold Spring Harbor Laboratory, Cold Spring Harbor, New York.
- [29] Rohlf, J.F. (1994) *NTSYS. Numerical Taxonomy and Multivariate Analysis System*. Exeter software, New York.
- [30] Maidak, B.L., Cole, J.R., Lilburn, T.G., Parker Jr., C.T., Saxman, P.R., Farris, R.J., Garrity, G.M., Olsen, G.J., Schmidt, T.M. and Tiedje, J.M. (2001) The RDP-II (Ribosomal Database Project). *Nucleic Acids Res.* 29, 173–174.
- [31] Altschul, S.F., Madden, T.L., Schäffer, A.A., Zhang, J., Zhang, Z., Miller, W. and Lipman, D.J. (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* 25, 3389–3402.
- [32] Pearson, W.R. and Lipman, D.J. (1988) Improved tools for biological sequence comparison. *Proc. Natl. Acad. Sci. USA* 85, 2444–2448.
- [33] Thompson, J.D., Gibson, T.J., Plewniak, F., Jeanmougin, F. and Higgins, D.G. (1997) The ClustalX windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.* 24, 4876–4882.
- [34] Felsenstein, J. (1993) *PHYLIP – phylogeny Inference Package Version 3.5c*. Department of Genetics, University of Washington, Seattle (distributed by the author).
- [35] Marilley, L. and Aragno, M. (1999) Phylogenetic diversity of bacterial communities from different proximity to *Lolium perenne* and *Trifolium repens* roots. *Appl. Soil Ecol.* 13, 127–136.
- [36] Boeckelmann, U., Manz, W., Neu, T.R. and Szewzyk, U. (2000) Characterization of the bacterial community of lotic organic aggregates ('river snow') in the Elbe River of Germany by cultivation and molecular methods. *FEMS Microbiol. Ecol.* 33, 157–170.
- [37] Orpin, C.G., Anderson, J.M. (1988) The animal environment. In: *Micro-Organisms in Action: Concepts and Applications in Microbial Ecology* (Lynch, J.M., Hobbie, E.J. Eds), pp. 136–192. Blackwell Scientific Publications, London.
- [38] Daniel, O. and Anderson, J.M. (1992) Microbial biomass and activity in contrasting soil materials after passage through the gut of the earthworm *Lumbricus rubellus*, Hofmeister. *Soil Biol. Biochem.* 18, 465–470.
- [39] Clegg, C.D., van Elsas, J.D., Anderson, J.M. and Lapin-Scott, H.M. (1994) Assessment of the role of a terrestrial isopod in the survival of a genetically modified pseudomonad and its detection using the polymerase chain reaction. *FEMS Microb. Ecol.* 15, 161–168.
- [40] Wood, S. and Griffiths, B.S. (1988) Bacteria associated with the hepatopancreas of the woodlice *Oniscus asellus* and *Porcellio scaber* (Crustacea, Isopoda). *Pedobiologia* 31, 89–94.
- [41] Hanlon, R.D.G. and Anderson, J.M. (1980) Influence of macroarthropod feeding activities on microflora in decomposing oak leaves. *Soil Biol. Biochem.* 12, 255–261.
- [42] Torsvik, V., Goksoyr, J. and Daae, F.L. (1990) High diversity in DNA of soil bacteria. *Appl. Environ. Microbiol.* 56, 782–787.
- [43] Hugenholtz, P., Goebel, B.M. and Pace, N.R. (1998) Impact of culture-independent studies on the emerging phylogenetic view of bacterial diversity. *J. Bacteriol.* 180, 4765–4774.
- [44] Sneath, P.H.A., Mair, N.S., Sharpe, M.E., Holt, J.G. (1984) *Bergey's Manual of Systematic Bacteriology*, Vol. 2. Williams and Wilkins, Baltimore, MD.
- [45] Stewart, C.S., Flint, H.J., Bryant, M.P. (1997) The rumen bacteria. In: *The Rumen Microbial Ecosystem* (Hobson, P.N., Stewart, C.S., Eds), pp. 10–72. Blackie Academic and Professional, London.
- [46] Mackie, R.I., White, B.A., Isaacson, R.E. (1997) *Gastrointestinal Microbiology*. Vol. 2. *Gastrointestinal Microbes and Host Interactions*. Chapman and Hall Microbiology Series, New York.