

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/329130913>

# Apparent kleptoparasitism in fish—parasitic gnathiid isopods

Article in *Parasitology Research* · November 2018

DOI: 10.1007/s00436-018-6152-8

CITATIONS

7

READS

147

4 authors:



**Mary Shodipo**

Silliman University

5 PUBLICATIONS 16 CITATIONS

[SEE PROFILE](#)



**Rickdane Gomez**

Marine Science Institute

2 PUBLICATIONS 7 CITATIONS

[SEE PROFILE](#)



**Rachel Welicky**

University of Washington Seattle

32 PUBLICATIONS 266 CITATIONS

[SEE PROFILE](#)



**Paul Sikkel**

University of Miami

110 PUBLICATIONS 1,597 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:




Gnathiid isopod host and habitat associations in the Philippines [View project](#)



Ectoparasite Ecology [View project](#)



# Apparent kleptoparasitism in fish—parasitic gnathiid isopods

Mary O. Shodipo<sup>1</sup> · Rick Dane C. Gomez<sup>2</sup> · Rachel L. Welicky<sup>3</sup> · Paul C. Sikkel<sup>3,4</sup> 

Received: 19 July 2018 / Accepted: 12 November 2018  
© Springer-Verlag GmbH Germany, part of Springer Nature 2018

## Abstract

Gnathiid isopods are common external parasites/micropredators that feed on the blood of marine fishes. During the course of processing samples of gnathiid isopods collected from light traps in the central Philippines, we observed a gnathiid attached to and apparently feeding from the abdomen of another gnathiid. Because the abdomens of both gnathiids were enlarged, it was unclear whether one actually fed on the blood meal of the other. Introduction of unfed gnathiids with fed gnathiids revealed that one gnathiid could and did feed on the blood meal of another. This is the first observation of apparent conspecific kleptoparasitism reported for gnathiid isopods.

**Keywords** Micropredator · Hematophagous · Coral reef · Philippines

## Introduction

Parasitism is one of the most common consumer strategies (e.g., Weinstein and Kuris 2016) and the duration of infection ranges from acute (one-time) to chronic and permanent (facultative obligate). Parasites that spend much or most of their life history as free-living and only associate temporarily with hosts may more appropriately be considered “micropredators” (Kuris and Lafferty 2000; Lafferty and Kuris 2002).

Perhaps the best-studied micropredators are blood-feeding arthropods such as mosquitoes and ticks (e.g., Mans and Nietz 2004; Gargili et al. 2017). These organisms attach temporarily while feeding on the blood of vertebrate hosts. The uptake of blood from a host typically occurs from the insertion of a

proboscis into a blood vessel of a host, and the blood meal is clearly visible through the translucent abdomen.

Gnathiid isopods (Isopoda: Gnathiidae) are the marine equivalent of terrestrial ticks and mosquitoes. They have three larval phases (Smit and Davies 2004; Tanaka 2007) and to molt from one phase to the next they must obtain a blood meal from a fish host. Unfed gnathiids are considered “zuphea” and following each blood meal, they become “pranizae” with a clearly distended abdomen. After the third blood meal, they metamorphose into sexually dimorphic non-parasitic adults (Smit and Davies 2004; Tanaka 2007). Gnathiids infest a wide range of hosts to obtain their blood meals (Ferreira et al. 2009; Coile and Sikkel 2013), and accordingly are referred to as generalists with preferences (Jones et al. 2007; Nagel and Grutter 2007).

Given that the blood meal obtained by blood-feeding arthropods is clearly visible through the abdomen and that the thin abdominal tissue likely allows for the passage of chemical cues, it seems possible that blood-feeding arthropods could feed on the blood meals of one-another. Yet, to our knowledge, this has never been reported.

## Methods, results, and discussion

As part of a continuing study on gnathiid isopod host and habitat associations (Sikkel et al. 2014, Santos and Sikkel 2017), in 2018 on Siquijor, central Philippines (9° 8'39.30" N, 123°30'29.30"E), we collected gnathiids using light traps (Artim and Sikkel 2016). After the light traps were collected,

---

Section Editor: Simonetta Mattiucci

✉ Paul C. Sikkel  
paul.sikkel@gmail.com

<sup>1</sup> Institute of Environmental and Marine Sciences, Silliman University, 6200 Dumaguete City, Negros Oriental, Philippines

<sup>2</sup> Department of Biology, Ateneo de Manila University, Katipunan Ave., 1105 Quezon City, Philippines

<sup>3</sup> Water Research Group, Unit for Environmental Sciences and Management, North-West University, Potchefstroom 2520, South Africa

<sup>4</sup> Department of Biology, Arkansas State University, PO Box 599, State University, AR 72467, USA

the benthic fauna from the light trap was surveyed for gnathiids under stereomicroscopes. While searching for gnathiids, in 8 instances, three researchers (PCS, MOS, and RCG) independently observed two gnathiids that remained still and connected to one another. Both gnathiids had distended abdomens, but the ventral side of the cephalon of one gnathiid, where the mouthparts are situated, were laid directly onto the pereon of the other, where the blood meal is located. This suggested the possibility that one zuphea gnathiid was obtaining a blood meal from a praniza gnathiid. Based on these observations alone, it was not possible to determine if this was the case. Thus, we introduced 30 zuphea in a 10-cm diameter plastic dish with 6 pranizae. Sample sizes were opportunistic and based on the number of live gnathiids that were collected and available at the time. After 3 h, one zuphea attached to one praniza and the pair was monitored continuously for 1.5 h during which they remained attached to one another. Observations discontinued for 8.5 h during the night, but the pair was in contact the following morning and remained so for an additional 2 h before detaching. During attachment, the feeding structure of the zuphea was observed to be inserted into the pereon of the praniza and the abdomen of the zuphea became clearly distended, while the abdomen of the praniza became less so (Fig. 1). For the gnathiid that had its blood meal “stolen,” we did not observe any immediate negative effects such as injury or death. In a second treatment, 12 dead pranizae and 9 live zuphea were placed together in a plastic dish over 13 h, and none of the zuphea attached to the dead pranizae.

Our observations suggest that ectoparasitic gnathiid isopods can obtain some blood meals from fish hosts indirectly. There are two possible explanations for indirect feeding. First, indirect feeding may simply be a bi-product of the use of chemical cues to locate fish hosts, combined with the close proximity of gnathiids to one another in dishes. Although



**Fig. 1** Fed gnathiid isopod (praniza) having its blood meal taken directly by an unfed gnathiid (zuphea)

gnathiid host-finding mechanisms are poorly understood, studies have shown that gnathiids appear to be able to use chemical cues, or a combination of visual and chemical cues to find hosts (Nagel et al. 2008; Sikkell et al. 2011; Jenkins et al. 2018). Chemical cues may permeate through the abdomen of a recently fed praniza and be detectable by a zuphea, thereby attracting it to a “host.” This postulate is supported by findings from mosquitoes that integrate use of thermal, olfactory, and visual cues to locate hosts (reviewed in Gibson and Torr 1999; McMeniman et al. 2014; Van Breugel et al. 2015). If such a multi-sensory strategy is also utilized by gnathiids, then it may explain why no zuphea attached and fed on dead pranizae in the second treatment. The chemical cues produced by decomposing blood from a dead gnathiid are also likely different from those produced by freshly ingested blood.

A second, and non-mutually exclusive, explanation for this observed feeding behavior is that it actually reflects a kleptoparasitic strategy. Kleptoparasitism is the act of one organism stealing food from another, and these interactions are most commonly described from terrestrial organisms. Nevertheless, kleptoparasitism does occur in aquatic systems, with the majority of examples involving interactions between invertebrates (reviewed in Iyengar 2008). For example, marine snails such as *Trichotropis cancellata* can kleptoparasitize tube polychaete worms by stealing food directly from their mouth using their pseudoproboscis (Iyengar 2002). *Trichotropis cancellata* are also suspension feeders so are facultative kleptoparasites, however kleptoparasitism appears to provide a growth advantage, as snails using only this strategy were found to grow quicker than those that were purely suspension feeders, albeit at the expense of their host who’s growth rates were reduced (Iyengar 2002, 2004).

Marine organisms can also be obligate kleptoparasites; however, this occurs less frequently (reviewed by Iyengar 2008). Maine copepods *Ascidicola rosea* appear to obtain their food solely from the sea squirt *Corella parallelogramma*, it positions itself in the oesophageal region and feeds directly from the host’s food (Gotto 1957). *Corella parallelogramma* can host as many as 8 different copepod species this way (Gotto 1957). Intraspecific marine kleptoparasitic interactions appear less common, and of those reported, these interactions seemingly represent cooperative hunting and social foraging activity patterns rather than true kleptoparasitism (i.e., Vijai et al. 2017).

For gnathiids living in high densities or where hosts are rare, kleptoparasitism may reduce the energetic costs associated with locating a susceptible host to infest. However, though no immediate negative effects were observed for the gnathiid that was fed upon, potentially the loss of its blood meal may delay or prevent its further development. Although we are unaware of the frequency with which this interaction occurs, the reduced necessity for some gnathiids to find a fish host to infest for obtaining a blood

meal provides insight into how gnathiid interactions may change if fish host populations decline.

**Acknowledgements** We thank the municipality of San Juan, Siquijor, Philippines, for permission to conduct this study. We also thank Dr. Hilconida P. Calumpong, Dr. Janet S. Estacion, Dr. Rene A. Abesamis, and the staff of the Silliman University Institute for Environmental and Marine Sciences for logistic support.

**Funding information** Funding was generously provided by grants from the National Geographic Society and the US National Science Foundation (OCE-1536794, PC Sikkell, PI). Financial assistance for RL Welicky from the Claude Leon Foundation of South Africa for this research is also acknowledged. This is contribution number 268 from the NWU-Water Research Group.

### Compliance with ethical standards

**Ethical approval** All applicable international, national, and/or institutional guidelines for the care and use of animals were followed. All procedures performed in studies involving animals were in accordance with the ethical standards of Silliman University, Arkansas State University, North-West University, and the Government of the Philippines.

**Conflict of interest** The authors declare that they have no conflict of interest.

### References

- Artim JM, Sikkell PC (2016) Comparison of sampling methodologies and estimation of population parameters for a temporary fish ectoparasite. *Int J Parasitol Parasites Wildl* 5:145–157
- Coile AM, Sikkell PC (2013) An experimental field test of susceptibility to ectoparasitic gnathiid isopods among Caribbean reef fishes. *Parasitology* 140:888–896
- Ferreira ML, Smit NJ, Grutter AS, Davies AJ (2009) A new species of gnathiid (Crustacea: Isopoda) parasitizing teleosts from Lizard Island, Great Barrier Reef, Australia. *J Parasitol* 95:1066–1075
- Gargili A, Estrada-Pena A, Spengler JR, Lukashev A, Nuttall PA, Bente DA (2017) The role of ticks in the maintenance and transmission of Crimean-Congo hemorrhagic fever virus: a review of published field and laboratory studies. *Antivir Res* 144:93–119
- Gibson G, Torr SJ (1999) Visual and olfactory responses of haematophagous Diptera to host stimuli. *Med Vet Entomol* 13:2–23
- Gotto RV (1957) The biology of a commensal copepod, *Ascidicola rosea* Thorell, in the ascidian *Corella parallelogramma* (Müller). *J Mar Biol Assoc U K* 36:281–290
- Iyengar EV (2002) Sneaky snails and wasted worms: kleptoparasitism by *Trichotropis cancellata* (Mollusca, Gastropoda) on *Serpula columbiana* (Annelida, Polychaeta). *Mar Ecol Prog Ser* 244:153–162
- Iyengar EV (2004) Host-specific performance and host use in the kleptoparasitic marine snail *Trichotropis cancellata*. *Oecologia* 138:628–639
- Iyengar EV (2008) Kleptoparasitic interactions throughout the animal kingdom and a re-evaluation, based on participant mobility, of the conditions promoting the evolution of kleptoparasitism. *Biol J Linn Soc* 93:745–762
- Jenkins WG, Demopoulos AW, Sikkell PC (2018) Effects of host injury on susceptibility of marine reef fishes to ectoparasitic gnathiid isopods, vol 75. *Symbiosis*, pp 113–121
- Jones CM, Nagel L, Huges GL, Cribb TH, Grutter AS (2007) Host specificity of two species of *Gnathia* (Isopoda) determined by DNA sequencing blood meals. *Int J Parasitol* 37:927–935
- Kuris AM, Lafferty KD (2000) Parasite-host modeling meets reality: adaptive peaks and their ecological attributes. In: Poulin R, Morand S, Skorping A (eds) *Evolutionary biology of host-parasite relationships: theory meets reality*. Elsevier Science, New York, pp 9–26
- Lafferty KD, Kuris AM (2002) Trophic strategies, animal diversity and body size. *Trends Ecol Evol* 17(11):507–513
- Mans BJ, Neitz AW (2004) Adaptation of ticks to a blood-feeding environment: evolution from a functional perspective. *Insect Biochem Mol Biol* 34(1):1–17
- McMeniman CJ, Corfas RA, Matthews BJ, Ritchie SA, Vossshall LB (2014) Multimodal integration of carbon dioxide and other sensory cues drives mosquito attraction to humans. *Cell* 156(5):1060–1071
- Nagel L, Grutter AS (2007) Host preference and specialization in *Gnathia* sp., a common parasitic isopod of coral reef fishes. *J Fish Biol* 70:497–508
- Nagel L, Montgomerie R, Lougheed S (2008) Evolutionary divergence in common marine ectoparasites *Gnathia* spp. (Isopoda: Gnathiidae) on the Great Barrier Reef: phylogeography, morphology, and behaviour. *Biol J Linn Soc* 94:569–587
- Santos TRN, Sikkell PC (2017) Habitat associations of fish-parasitic gnathiid isopods in a shallow reef system in the central Philippines. *Mar Biodivers* 1–14
- Sikkell PC, Sears WT, Weldon B, Tuttle BC (2011) An experimental field test of host-finding mechanisms in a Caribbean gnathiid isopod. *Mar Biol* 158:1075–1083
- Sikkell PC, Tuttle LJ, Cure K, Coile AM, Hixon MA (2014) Low susceptibility of invasive red lionfish (*Pterois volitans*) to a generalist ectoparasite in both its introduced and native ranges. *PLoS One* 9(5):e95854
- Smit NJ, Davies AJ (2004) The curious life-style of the parasitic stages of gnathiid isopods. *Adv Parasitol* 58:289–391
- Tanaka K (2007) Life history of gnathiid isopods-current knowledge and future directions. *Plankton Benthos Res* 2:1–11
- Van Breugel F, Riffell J, Fairhall A, Dickinson MH (2015) Mosquitoes use vision to associate odor plumes with thermal targets. *Curr Biol* 25:2123–2129
- Vijai D, Puneeta P, Matsui H, Morooka M, Sakurai Y (2017) Intraspecific kleptoparasitism in squid school and schooling behavior of its prey. *Mar Biodivers* 47:637–638
- Weinstein SB, Kuris AM (2016) Independent origins of parasitism in Animalia. *Biol Lett* 12:20160324